



BMJ Open is committed to open peer review. As part of this commitment we make the peer review history of every article we publish publicly available.

When an article is published we post the peer reviewers' comments and the authors' responses online. We also post the versions of the paper that were used during peer review. These are the versions that the peer review comments apply to.

The versions of the paper that follow are the versions that were submitted during the peer review process. They are not the versions of record or the final published versions. They should not be cited or distributed as the published version of this manuscript.

BMJ Open is an open access journal and the full, final, typeset and author-corrected version of record of the manuscript is available on our site with no access controls, subscription charges or pay-per-view fees (<http://bmjopen.bmj.com>).

If you have any questions on BMJ Open's open peer review process please email [info.bmjopen@bmj.com](mailto:info.bmjopen@bmj.com)

# BMJ Open

## **Insights into the impact of ACEI/ARBs on COVID-19 prognosis: A multi-state model of nationwide hospital surveillance data**

Journal:	<i>BMJ Open</i>
Manuscript ID	bmjopen-2021-053393
Article Type:	Original research
Date Submitted by the Author:	13-May-2021
Complete List of Authors:	Penalvo, Jose; Institute of Tropical Medicine Genbrugge, Els; Institute of Tropical Medicine Mertens, Elly; Institute of Tropical Medicine, Sagastume, Diana; Institute of Tropical Medicine van der Sande, Marianne A. B.; Institute of Tropical Medicine Widdowson, Marc-Alain; Institute of Tropical Medicine Van Beckhoven, Dominique; Sciensano
Keywords:	COVID-19, Hypertension < CARDIOLOGY, STATISTICS & RESEARCH METHODS

SCHOLARONE™  
Manuscripts



I, the Submitting Author has the right to grant and does grant on behalf of all authors of the Work (as defined in the below author licence), an exclusive licence and/or a non-exclusive licence for contributions from authors who are: i) UK Crown employees; ii) where BMJ has agreed a CC-BY licence shall apply, and/or iii) in accordance with the terms applicable for US Federal Government officers or employees acting as part of their official duties; on a worldwide, perpetual, irrevocable, royalty-free basis to BMJ Publishing Group Ltd ("BMJ") its licensees and where the relevant Journal is co-owned by BMJ to the co-owners of the Journal, to publish the Work in this journal and any other BMJ products and to exploit all rights, as set out in our [licence](#).

The Submitting Author accepts and understands that any supply made under these terms is made by BMJ to the Submitting Author unless you are acting as an employee on behalf of your employer or a postgraduate student of an affiliated institution which is paying any applicable article publishing charge ("APC") for Open Access articles. Where the Submitting Author wishes to make the Work available on an Open Access basis (and intends to pay the relevant APC), the terms of reuse of such Open Access shall be governed by a Creative Commons licence – details of these licences and which [Creative Commons](#) licence will apply to this Work are set out in our licence referred to above.

Other than as permitted in any relevant BMJ Author's Self Archiving Policies, I confirm this Work has not been accepted for publication elsewhere, is not being considered for publication elsewhere and does not duplicate material already published. I confirm all authors consent to publication of this Work and authorise the granting of this licence.

**Insights into the impact of ACEI/ARBs on COVID-19 prognosis: A multi-state model of nationwide hospital surveillance data**

José L. Peñalvo<sup>1</sup>, Els Genbrugge<sup>1</sup>, Elly Mertens<sup>1</sup>, Diana Sagastume<sup>1</sup>, Marianne A. B. van der Sande<sup>2,3</sup>, Marc-Alain Widdowson<sup>4</sup>, Dominique Van Beckhoven<sup>5</sup>

<sup>1</sup> Non-Communicable Diseases Unit, Department of Public Health, Institute of Tropical Medicine, Antwerp, Belgium

<sup>2</sup> Department of Public Health, Institute of Tropical Medicine, Antwerp, Belgium

<sup>3</sup> Global Health Julius Centre for Health Sciences and Primary Care, University Medical Centre Utrecht, Utrecht, The Netherlands.

<sup>4</sup> Institute of Tropical Medicine, Antwerp, Belgium

<sup>5</sup> Department of Epidemiology and Public Health, Sciensano, Brussels, Belgium

**Correspondence**

José L. Peñalvo, Non-Communicable Diseases Unit, Department of Public Health, Institute of Tropical Medicine. Nationalestraat 150, 2000-Antwerp, Belgium. Tel: +32(0)32476251, email [jpenalvo@itg.be](mailto:jpenalvo@itg.be)

**Keywords**

Angiotensin-converting enzyme inhibitors (ACEI); Angiotensin-receptor blockers (ARBs); COVID-19; comorbidities; multi-state model.

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

**Abstract**

*Objectives:* The widespread use of angiotensin-converting enzyme inhibitors (ACEI) and angiotensin receptor blockers (ARB) in patients by chronic patients raised early concerns on the potential exacerbation of COVID-19 severity and fatality. A number of conflicting studies have used standard methods that may lead to biased estimates when analyzing hospital data because of the presence of competing events and time-dependent complexity. We investigated the effect of ACEI/ARBs use COVID-19 disease outcomes using time-to-event data in a multi-state setting to account for competing events and minimize bias.

*Setting:* Nationwide surveillance data from 119 Belgian hospitals.

*Participants:* Medical records of 10,866 patients hospitalised from March 14 to June 14, 2020 with a confirmed SARS-CoV-19 infection and information about ACEI/ARBs use.

*Primary outcome measure:* Multi-state, multivariate Cox-Markov models were used to estimate the hazards of patients transitioning through health states from admission to discharge or death, along with transition probabilities calculated by combining the baseline cumulative hazard and regression coefficients.

*Results:* After accounting for potential confounders no evidence was found of a detrimental effect of ACEI/ARBs use on admission to intensive care (ICU) or on in-hospital death. Contrastingly, ACEI/ARBs use was associated with a modest positive effect on recovery (HR 1.07 [95%CI 1.01-1.13],  $p=0.027$ ) and reduced fatality (0.83, 0.75-0.93,  $p=0.001$ ). For patients needing ICU admission, no evidence of an association between ACEI/ARBs use and recovery (1.16, 0.97-1.38,  $p=0.098$ ) or in-hospital death during ICU (0.91, 0.73-1.12,  $p=0.381$ ) was observed. Male gender and older age were significantly associated with higher risk of ICU admission or death. Chronic cardiometabolic comorbidities were also associated with less recovery.

*Conclusions:* For the first time, a multistate model was used to address magnitude and direction of the effect of ACEI/ARBs use on COVID-19 progression. By minimizing bias, this study provided robust indication a protective, albeit modest, effect on recovery and survival.

**Keywords:** Angiotensin-converting enzyme inhibitors (ACEI); Angiotensin-receptor blockers (ARBs); COVID-19; comorbidities; multi-state model.

### **Strengths and limitations of this study**

- The study uses nationwide hospital surveillance data, and includes all general hospitals in Belgium.
- The use of a comprehensive database, but more so the utilization of models that adequately fitting to time-to-event hospital data with mutually exclusive health states results in less probability of introducing biases and are crucial for correct evidence-based information for decision making.
- Only transfer to intensive care was linked to a calendar date and was therefore the only event which could be used as a proxy for severe disease state in our time-dependent model, indicating that our estimates might represent more a critical state of the patient.
- Our analysis provided robust estimation a protective effect on ACEI/ARBs in recovery and survival of hospitalized COVID-19 patients.

1. BACKGROUND

COVID-19 virus infection is likely to cause severe disease among older individuals, men and patients with chronic respiratory or cardiometabolic conditions such as cardiovascular disease (CVD), hypertension (HTN) and diabetes (DM) <sup>1-3</sup>. Also, common risk factors for chronic conditions, such as smoking and particularly obesity, have been identified as key predictors of hospitalization and critical illness, even in young adults with no underlying conditions <sup>4,5</sup>. While the pathogenesis of certain chronic diseases predisposes to severe COVID-19 outcomes, common medications might also increase this risk because of the interaction between SARS-CoV-2 and the angiotensin-converting enzyme 2 (ACE2)<sup>6</sup>, an enzyme that physiologically counters the renin-angiotensin-aldosterone system (RAAS) activation.

SARS-CoV-2 binds to target cells using the angiotensin-converting enzyme 2 (ACE2) in cell membranes <sup>6</sup>, a component of the RAAS, that can degrade angiotensin II to attenuate its subsequent physiological action (vasoconstriction). Modulation of RAAS is a frequent mode of action of antihypertensive and hypoglycemic drugs. HTN is commonly treated with ACE inhibitors (ACEI) and angiotensin II receptor blockers (ARBs) which results in an upregulation of ACE2. Expression of ACE2 is substantially increased in patients with HTN or type 1 or 2 DM, who are treated with ACEI or angiotensin receptor blockers (ARBs), and this may explain the higher risk for severe or fatal covid-19 infection among this patients. This information suggests that ACE2 expression is increased in hypertensive and diabetic patients treated with ACEI and ARBs (and potentially other commonly used drugs for the management of chronic conditions) worsening the prognosis of COVID-19 infection. This raised immediate concerns during the first phases of the outbreak of SARS-CoV-2 pandemic because of the widespread used RAAS inhibitors, specifically ACEI or ARBs among chronic, mostly hypertensive or diabetic, patients <sup>7</sup>. As patients with chronic comorbidities were also identified as more vulnerable to severe COVID-19 disease, it is necessary to understand whether part of this vulnerability could be

1  
2  
3 96 attributed to the use of ACEI/ARBs and to evaluate the risk of discontinuing this otherwise essential,  
4  
5 97 first-line therapy, for hypertensive and diabetic patients.  
6

7 98 To date, a number of studies addressing the potential effect of ACEI and ARBs on the prognosis of  
8  
9 99 COVID-19 have been reported, mostly supporting the absence of harmful effects of these drugs on  
10  
11 100 COVID-19 prognosis<sup>8-10</sup>. In these studies, a wide range of statistical methods have been used to test  
12  
13 101 this hypothesis, including comparison of proportions, percentage points, logistic regression, or time-  
14  
15 102 to-event analysis and Cox models. The use of standard methods for these particular analyses can easily  
16  
17 103 lead to biased estimates, in particular when analyzing hospital data because of the presence of  
18  
19 104 competing events, such as death and recovery, and the time-dependency of these competing events  
20  
21 105<sup>11</sup>. As such, the analysis of the association of ACEI/ARBs on the progression of COVID-19 or related  
22  
23 106 mortality requires the assessment of competing risks/events. Analyzing time-to-event data in a multi-  
24  
25 107 state setting would better fit the true progression of COVID-19 in hospitalized patients, as shown by  
26  
27 108 two studies using the multi-state-approach in a COVID-19 context<sup>12,13</sup>. Multi-state models allow for  
28  
29 109 studying clinically competing events (discharged alive vs deceased), as well as disease progression (in  
30  
31 110 terms of hospital stay duration, transfer to Intensive Care Unit (ICU), treatment received),  
32  
33 111 simultaneously over time. This multi-state model framework ensures avoiding bias that stems from  
34  
35 112 censoring patients (informative censoring bias), as well as circumvents time-dependent bias by  
36  
37 113 treating disease progression as a transient state that might influence the probability of experiencing a  
38  
39 114 certain future outcome depending on patient's risk factors. While accounting for these biases, we  
40  
41 115 revisited the the hypothesis of the potential effect of ACEI/ARBs use in patient's prognosis during  
42  
43 116 hospitalization using a competing risk multi-state model and nationwide hospital surveillance data on  
44  
45 117 COVID-19 patients in Belgium.  
46  
47  
48  
49  
50  
51  
52  
53

## 54 119 **2. PATIENTS AND METHODS**

### 55 120 **2.1. Data sources**

56  
57  
58  
59  
60



1  
2  
3 121 All methods were carried out in accordance with relevant guidelines and regulations. Nationwide  
4  
5 122 hospital surveillance data on COVID-19 patients in Belgium are routinely gathered by Sciensano, the  
6  
7 123 Belgian Institute of Public Health, which is legally entitled institution for surveillance of infectious  
8  
9 124 diseases in Belgium (Royal Decree of 21/03/2018). Retrieving informed consent was determined as a  
10  
11 125 disproportional load on the hospital resources in the crisis situation. An information letter was given  
12  
13 126 to the patients at the time of discharge which contained an explanation of their rights concerning the  
14  
15 127 data that was gathered by Sciensano. The COVID-19 hospital surveillance was authorised by an  
16  
17 128 independent administrative authority protecting privacy and personal data and was approved by the  
18  
19 129 ethical committee of Ghent University Hospital (BC-07507). Details on the Belgian COVID-19 hospital  
20  
21 130 surveillance system have been previously published <sup>14</sup>. The system cover 119 hospitals in Belgium, who  
22  
23 131 report standardized information on hospitalized COVID-19 patients collected through a structured  
24  
25 132 questionnaire at hospital admission and discharge. An anonymized subset of data from Sciensano was  
26  
27 133 shared with the Institute of Tropical Medicine through a secured data transfer platform applying data  
28  
29 134 encryption. Ethical clearance was obtained from the Institutional Review Board (IRB) of the Institute  
30  
31 135 of Tropical Medicine after revision of the research protocol num. 1393/20, 02/05/2020.  
32  
33  
34 136 Variables collected at admission include the date of hospital admission, reason for hospitalization,  
35  
36 137 symptoms, clinical signs, treatment with ACEI or ARBs, and demographic information such as age,  
37  
38 138 gender, and the presence of chronic comorbidities. Information recorded at discharge includes  
39  
40 139 laboratory values, details on COVID-19 specific treatments during hospital stay, date of discharge,  
41  
42 140 health status at discharge, and measures on the severity of the disease such as the need for transfer  
43  
44 141 to ICU, invasive ventilation support and/or oxygenation by extracorporeal membrane oxygenation  
45  
46 142 (ECMO), and the development of a bacterial and/or fungal superinfection, pneumonia, and/or acute  
47  
48 143 respiratory distress syndrome (ARDS). Dates for these severe events were only available for ICU  
49  
50 144 transfer.  
51  
52 145  
53  
54 146  
55  
56  
57  
58  
59  
60

## 2.2. Study population

Adult COVID-19 patients with a SARS-CoV-19 infection confirmed by polymerase chain reaction (PCR), and/or suggestive imaging alterations on chest CT combined with typical clinical presentation, at admission or while hospitalized in Belgium from March 14 to June 14, 2020 were considered as COVID-19 patients (n=16,341). Of these, patients with completed questionnaires both on admission and discharge (12,109 patients, 74.1%) were selected. Information on patients admitted to hospitals before March 1 2020 (270, 1.65%) for reasons other than COVID-19 and infected while hospitalized was also removed. Furthermore, patients with implausible admission dates were removed, including: Date of discharge before date of admission (42 patients, 0.25%); Date of ICU transfer preceding date of admission (31 patients, 0.19%); Date of discharge before date of ICU transfer (2 patients, 0.01%); Date of discharge preceding the date of ICU discharge if the difference was more than 1 day (47 patients, 0.28%). The final dataset for descriptive analyses included information on 11717 COVID-19 patients. For the multivariate multi-state model, patients with unknown use of ACE/ARBs (718 patients, 6.12%) were also excluded, along with those missing information on gender (118 patients, 0.72%) or unknown transfer to ICU (15, 0.09%). The final dataset for the multi-state model contained information from 103 hospitals in Belgium and 10,866 COVID-19 patients, including 539 patients (5%) that were admitted directly to ICU.

## 2.3. Study outcomes

Patients were considered to have recovered when their status at discharge was recorded as “cured” or “other”. In the latter case, it was assumed they were allowed to recover at home, revalidation center or nursing home. Patients were considered to have an in-hospital death when their status at discharge was recorded as “death”. Patients were considered lost to follow up when their status at discharge was recorded as “unknown” or when they were transferred to another hospital (status at discharge = “transfer to another hospital”), as no further information was available. Severe COVID-19 was captured in the database as an illness that: required ECMO, or artificial ventilation, experienced ARDS, pneumonia, bacterial and/or fungal co-infection, or required transfer to or

1  
2  
3 173 treatment at ICU. Among these, event date was only available for transfer to ICU, and only this variable  
4  
5 174 could therefore be selected for the models as time-defined severity outcome. Based on this, time to  
6  
7 175 severe illness was defined as the time passed from hospital admission to date of transfer to ICU. Length  
8  
9  
10 176 of hospital stay was defined as date from hospital admission to the date of hospital discharge (either  
11  
12 177 recovery, in-hospital death, or lost to follow-up).

13  
14 178 **2.4. Information on ACEI/ARBs and conditions related to COVID-19 prognosis**

15  
16 179 A dedicated section in the admission dataset covered the use of ACEI, ARBs or both, without  
17  
18  
19 180 specification of the specific drug. The admission database contained information on the following  
20  
21 181 factors associated with COVID-19<sup>15</sup>. Demographics (age and gender), risk factors (current smoking,  
22  
23 182 high blood pressure (HBP), and obesity), prevalent comorbidities (DM, chronic renal disease, CVD,  
24  
25 183 chronic lung disease, chronic liver disease, malignant solid neoplasms, hematological cancers, and  
26  
27 184 immunosuppression). Smoking status was only available for 53% of the patients. Obesity presented a  
28  
29  
30 185 large number of missing values (33.2%) because this variable was only recorded after April 3, 2020.  
31  
32 186 Similarly, there was also number of missing values for cognitive issues (5.7%) as this variable was only  
33  
34 187 recorded after March 23, 2020.

35  
36 188 **2.5. Patient and public involvement**

37  
38  
39 189 As a secondary data analysis of COVID-19 surveillance data this study did not involve patients or  
40  
41 190 the public in the design, conduct or dissemination plans.

42  
43 191  
44  
45 192 **3. STATISTICAL MODEL**

46  
47 193 Patient's characteristics at admission, during ICU, and discharge were visualized on histograms and  
48  
49 194 summarized as means and standard deviation and counts and percentage for continuous and  
50  
51 195 categorical variables, respectively. Descriptive analyses were provided for patients overall and  
52  
53 196 stratified by ACEI/ARBs use, including unknown use.

54  
55  
56 197 To study the impact of ACEI/ARBs on COVID-19 progression on a multivariate multi-state model, a  
57  
58 198 first model for identifying confounders was used. A backwards stepwise logistic regression with  
59  
60

ACEI/ARBs use as dependent variable and including factors and conditions previously associated with COVID-19 outcomes present in the database<sup>16,17</sup>, was used to inform the selection of potential confounders based on Akaike's information criterion (AIC). The variables used in the variable selection model were: gender, age, HBP, CVD, DM, obesity, chronic renal disease, chronic liver disease, chronic lung disease, solid malignant neoplasms, hematological cancers, immunodepression, and cognitive impairment. Two models were used depending on the availability of data, a first model (model 1) excluding variables collected at a later date (obesity, and cognitive issues) and using a full dataset (N=10,866), and a complete case analysis (model 2) excluding patients with missing data for obesity and cognitive issues (N=7,303).

We devised a multi-state model reflecting the progression of patients from admission to discharge accounting for the patient's characteristics identified to be potential confounders, and introducing ACEI/ABRs use as dependent variable. The model starts with one initial state (hospitalization), a potential (transient) state defined as ICU transfer (as a proxy for severe COVID-19 disease, as only ICU transfer had an associated time in the database) and two absorbing states (in-hospital death, and recovery). The multi-state model is characterized by transition hazards between the states; defined as the instantaneous risk for moving from one state to another. The transitions hazards are used to calculate transition probabilities, as the conditional probabilities of experiencing future outcomes, given the history and a particular set of prognostic factors (model covariates) for a given patient. The four-state model thereby comprised five possible patient's transitions; 1) hospitalization to ICU, 2) hospitalization to recovery, 3) hospitalization to in-hospital death, 4) ICU to recovery, and 5) ICU to in-hospital death, as presented in [Figure 1](#). A Cox-Markov model for the regression on the transition specific hazards was fitted using the *coxph* and *msfit* functions in R *survival* package<sup>18</sup>. This approach is equivalent of constructing five separate Cox regression models, one for each transition hazard. The cumulative baseline transition hazard (all covariate values equal to the reference value) was estimated by the Breslow estimator with the Aalen estimator of variance<sup>19</sup>. We integrated these separate Cox models in a multi-state framework studying different outcomes simultaneously and allowing the

1  
2  
3 225 calculation of transition probabilities. The transition probabilities were then estimated by combining  
4  
5 226 the baseline cumulative hazard and regression coefficients. Using R, *mstate* package was used with  
6  
7 227 *msfit* function to obtain cumulative (baseline) transition hazards and the function *probtrans* to obtain  
8  
9 228 the transition probabilities<sup>20</sup>. Estimates obtained from the Cox-Markov models are displayed in a table  
10  
11  
12 229 and significance is established at the 5% significance level. Cumulative (baseline) transition hazard  
13  
14 230 plots and transition probability plots were also generated for visual aid. In a setting with covariates, a  
15  
16 231 regression model for the transition specific hazards was used, whereby the covariates may help to  
17  
18 232 explain the difference in transition hazards. Model diagnostics were performed to check model  
19  
20 233 assumptions of proportional hazards, linearity, and interactions. Assumptions to the Markov model  
21  
22  
23 234 were assessed by including time from hospital admission to ICU transfer in the model for transition 4  
24  
25 235 and transition 5. A relaxation of the Markov assumption was also explored in the analysis.  
26  
27  
28 236

29  
30 237 **4. RESULTS**

31  
32 238 **4.1. Descriptive analysis**

33  
34 239 From the 11,717 patients available for this analysis, almost of all them (94.2%) presented  
35  
36 240 symptoms or clinical signs compatible with COVID-19 at admission ([Supplemental Table 1](#)). Most  
37  
38 241 frequent symptoms were fever (61.3%) and cough (53.2%), and most frequent signs were abnormal  
39  
40 242 pulmonary imaging (63.1%) compatible with viral pneumonia, abnormal auscultation (44.8%), and  
41  
42 243 dyspnea (42.4%). On admission, 15.1% of patients had a record of taking ACEI, and 8.5% ARBs, with  
43  
44 244 only 0.4% taking both ACEI and ARBs. For the purpose of this analysis these patients were merged as  
45  
46 245 ACEI/ARBs users. The majority of patients (69.9%) were nonusers of ACEI/ARBs versus 23.9% of users,  
47  
48 246 and only for a small proportion (6.1%) of patients the use of ACEI/ARBs was unknown ([Table 1](#)). No  
49  
50 247 difference was seen in the frequency of signs and symptoms reported according to ACEI/ARBs use  
51  
52 248 ([Supplemental Table 1](#)). Patients using ACEI/ARBs were markedly older (median [IQR] age 76 [65-84]  
53  
54 249 years) than non-users (67 [53-81] while no sex-differences were observed. As expected, ACEI/ARBs  
55  
56 250 users presented more frequently (74.4%) HBP than non-users (39.2%), as well as chronic lung disease  
57  
58  
59  
60

(16.8% vs 14.4%), chronic renal disease (19.3% vs 11.1%), DM (33.3% vs 18.1%), and particularly CVD (53.1% vs 28.4%). Among ACEI/ARBs users with 15.1% of them suffering multiple comorbidities (HBP, DM and CVD) versus 4.9% in non-users ([Table 1](#)). During hospital stay, over 80% of COVID-19 patients suffered one severe episode of either pneumonia, superinfection, ARDS, or mechanical ventilation, and 25.9% of patients had two or more severe episodes ([Table 2](#)). The most common manifestation of COVID-19 severity was pneumonia (79.1%), followed by other infections (19.4%), ARDS (12.7%), and artificial ventilation (7.6%). Frequency of severe conditions was almost the same for both ACEI/ARBs users and non-users (27.8% versus 25.4%). Of all admitted patients, 1,518 (13.0%) were transferred to ICU, mostly those with severe pneumonia (93.7%), or in need of artificial ventilation (58.0%), and for a mean duration of  $11.5 \pm 10.7$  days. Transfer to ICU was marginally more frequent among ACEI/ARBs users (15.1% versus 12.1%). Almost 78% of the patients admitted to Belgian hospitals recovered from COVID-19, either in the hospital (51.2%) or at home or revalidation centre or nursing home (26.4%) after an average  $12.6 \pm 10.9$  days in the hospital. Only 2% was lost to follow-up (transferred to another health care provider or unknown status at discharge).

#### 4.2. Multi-state model

A multivariate state-arrival extended Cox-Markov model was used to study the potentially different progression of COVID-19 patients through health states during hospitalization according to the use of ACEI/ARBs. Possible transitions, and number of patients in each health state are represented in [Figure 1](#). The selection of variables for adjusting the models were based on backwards stepwise logistic regression of ACEI/ARBs use as a function of potential confounding factors associated with COVID-19 recorded at admission ([Table 2 Supplemental](#)). A first model (model 1) using all available patients, identified five variables associated with the use of ACEI/ARBs and COVID-19: male gender (OR 1.33, 95%CI 1.21-1.47), older age (1.01, 1.01- 1.02 per 1-year increase), and prevalent CVD (1.71, 1.55 -1.90), diabetes (1.38, 1.24-1.54), and HBP (5.65, 5.10-6.27). Additionally, a second model (model 2) was used in sensitivity analysis including additional covariates (prevalent obesity and cognitive impairment) that were only available in a subset (62%) of patients ([Table 2 Supplemental](#)). Because very few patients

1  
2  
3 277 were asymptomatic on admission it was deemed unnecessary to adjust the regression models for  
4  
5 278 severity of disease at admission. For 16 patients (0.1%) it was unknown whether they were transferred  
6  
7 279 to ICU. These patients are therefore excluded from the multistate Cox-Markov regression analysis.  
8  
9

10 280 Plots for the cumulative hazard and transition probability between health states considering  
11  
12 281 ACEI/ARBs use were obtained by setting all model covariates to reference values (female gender, no  
13  
14 282 CVD, no HBP, and no DM), and median (70 years) age [Figure 2](#). When looking at the cumulative hazard  
15  
16 283 for the five possible transitions ([Figure 2A](#)), the hazard for recovery was markedly greater than that of  
17  
18 284 in-hospital death. In comparison with the other cumulative hazards, the hazard for transfer to ICU was  
19  
20 285 substantially smaller, representative of most COVID-19 patients not needing intensive care, or not  
21  
22 286 meeting criteria for admission (for instance after evaluation of frailty, and chance of survival). Transfer  
23  
24 287 to ICU was associated with increased hazard for in-hospital death and reduced hazard for recovery.  
25  
26 288 The use of ACEI/ARBs was associated with a modest but significant effect on the hazard of transition 2  
27  
28 289 (more recovery) and 3 (less in-hospital death), from admission. The use of ACEI/ARBs was not observed  
29  
30 290 to be associated with transfer to ICU (transition 1), nor with recovery (transition 4) or in-hospital death  
31  
32 291 (transition 5) after ICU. Overall, the probability of being transferred to ICU was for most patients less  
33  
34 292 than that of recovery ([Figure 2B](#)). However, those needing ICU had a reduced probability of recovery  
35  
36 293 and greater probability to decrease in the hospital than those patients not transferred to ICU ([Figure](#)  
37  
38 294 [2C](#)).  
39  
40  
41  
42

43 295 The estimates for the transition hazards for ACEI/ARBs use accounting for identified  
44  
45 296 confounding in the potential association with COVID-19 severity/fatality are presented in [Table 3](#). In  
46  
47 297 multivariate models, the use of ACE/ARBs (HR 1.07, 95%CI 1.01-1.13) was associated with more  
48  
49 298 recovery, and less death (0.83, 0.75-0.93). Even though there was a significant association between  
50  
51 299 ACEI/ARBs use the hazard of more recovery (transition 2), and less in-hospital death (transition 3) this  
52  
53 300 effect is modest, especially when inspecting state-occupation probabilities ([Figure 1 Supplemental](#)). In  
54  
55 301 respect to the adjusting variables ([Table 3 Supplemental](#)), male gender and HBP were associated with  
56  
57 302 transfer to ICU (severity), and older age also influenced this transition. Similarly, male gender, and  
58  
59  
60



older age, as well as prevalent CVD, HBP or DM were associated with less recovery. Similar to transfer to ICU, fatal progression was associated with male gender and age, as well as with prevalent CVD. No other comorbidity included in this model (i.e. associated with ACEI/ARBs use) appeared to be associated with fatality. For severe patients (transferred to ICU) recovery or death depended mostly on age, albeit fatal COVID-19 associated with the presence of DM, and a lengthier period between admission and ICU were significant associated with less recovery after ICU ([Table 3 Supplemental](#)). The impact of further adjustment for variables identified during confounder selection (obesity and cognitive issues) in the state transition of COVID-19 patients during hospitalization, resulted in loss of more than half of all patients due to missing values ([Table 1](#)). While estimated hazards for previous factors remained similar, the presence of cognitive issues was statistically significantly associated with transitions 1, 2, and 3 (i.e. less transfer to ICU, less recovery, and more in-hospital death), and obesity was strongly and statistically significantly associated with transition 1 only (more transfer to ICU). In this complete-case model, after additional adjustment for obesity and cognitive issues, the HR for ACEI/ARBs use for transition 2 (admission to recovery) is not anymore significant probably due to a decreased statistical power, since the point estimates remained similar.

## 5. DISCUSSION

In this study, a competing risk multistate model has been developed for the first time to address the magnitude and direction of the effect of ACEI/ARBs use in COVID-19 prognosis. Our analyses indicate a protective effect of ACEI/ARBs use, with increase recovery and survival, once important confounding factors such as age, particularly 70 and over, and male gender are accounted for. Chronic comorbidities such as CVD, HBP, and DM are also associated with less recovery in this model setting. Although there is a protective effect of ACEI/ARBs use on COVID-19 in-hospital death and more recovery, this effect is modest, especially when looking at the state-occupation probabilities. In our model, once the patient progresses to a severe state, no effect of ACEI/ARBs use was observed in the transition probabilities to recovery or in-hospital death; only older age and prevalent DM,



1  
2  
3 329 remained significant covariates in our model, arguably because of the smaller sample size (transfer to  
4  
5 330 ICU occurred only for 13% of patients). Previous studies using the same data source identified other  
6  
7 331 comorbidities as independent risk factors for COVID-19 severity/death in ICU patients, including  
8  
9 332 chronic pulmonary disease, chronic renal disease, and immunosuppression <sup>21</sup>. Although we  
10  
11 333 accounted for these factors in our model selection, they were not selected as they are not considered  
12  
13 334 to be related to the use of ACEI/ARBs but may nonetheless be independent risk factors for severity.  
14  
15

16 335       Because of the high clinical relevance, there have been numerous reports on studies of the  
17  
18 336 potential association between ACEI/ARBs and (worse) prognosis of COVID-19. Early studies of smaller  
19  
20 337 sample size and mostly descriptive design pointed to either no association or moderately lower rates  
21  
22 338 of severe disease among ACEI/ARBs users <sup>22-27</sup>. Further retrospective analysis involving larger patient  
23  
24 339 samples generally reported a lack of association <sup>8</sup>. A population-based study in Italy's Lombardy region  
25  
26 340 involving 6,272 cases identified across the Regional Health Service and matched 1:5 to population-  
27  
28 341 based control, found no association between the use of ACEI (adjusted OR 0.91, [95%CI 0.69-1.21]) or  
29  
30 342 ARBs (adjusted OR 0.83, [95%CI 0.63-1.10]) and severe/fatal COVID-19 <sup>28</sup>. Similarly, a case-control  
31  
32 343 study in the Spanish region of Madrid with data on 1,139 hospitalized cases matched 1:10 to  
33  
34 344 population controls found no association (adjusted OR 0.94, [95%CI 0.77-1.15]) of ACEI/ARBs use and  
35  
36 345 severe or fatal disease <sup>29</sup>. Analyzing data from all patients in the New York University Langone Health  
37  
38 346 electronic health record who had COVID-19 test results (12,594 patients), neither increased likelihood  
39  
40 347 of a positive test nor severe disease was observed for patients using ACEI/ARBs (or any other RAAS  
41  
42 348 medication) using propensity score matching <sup>30</sup>. In a nationwide study in Korea using claim records of  
43  
44 349 66,793 individuals tested for COVID-19, the use of ACEI/ARBs was not associated with a higher risk of  
45  
46 350 mortality (adjusted OR 0.88, [95%CI 0.53-1.44]) <sup>31</sup>. Similarly, a large retrospective analysis of an Italian  
47  
48 351 registry cohort including 43,000 patients concluded that neither ACEI (adjusted HR 0.97, [95%CI 0.89-  
49  
50 352 1.06]) or ARB (adjusted HR 0.98, [95% CI 0.89-1.06]) use was associated with either an increased or  
51  
52 353 decreased risk of all-cause mortality <sup>32</sup>. A multicenter study with 1,128 hypertensive patients, and using  
53  
54 354 mixed-effect Cox models (site as a random effect, and model adjusted for age, gender, comorbidities,  
55  
56  
57  
58  
59  
60

and in-hospital medications) reported a lower risk for all-cause mortality in the ACEI/ARB patients versus the non-ACEI/ARB group (adjusted HR 0.42, [95%CI 0.19-0.92]), and further compared with the use of other antihypertensive drugs, (adjusted HR 0.30, [95%CI, 0.12-0.70])<sup>33</sup>. Previous studies using Cox models reported also a reduced mortality risk for patients using ACEI/ARBs<sup>33,34</sup>. In others, albeit not statistically significant, estimates were very similar to the ones reported in our study for mortality (adjusted HR 0.83, [95%CI 0.67-1.03]) and for severe disease (adjusted HR 1.15, [95%CI 0.95-1.41])<sup>35</sup>. Similarly, but outside of the hospital setting, studies with data from general practitioners in England, found a strong association of ACEI/ARBs use and a reduced risk of COVID-19 disease (HR 0.63, [95%CI 0.59-0.67]) albeit not severity (HR 1.02, [95%CI 0.83-1.25]), and marked interactions with ethnicity with higher risks observed for Black Africans compared to Whites<sup>36</sup>. Variations between different ethnic groups raise the possibility of specific effects of ACEI/ARBs on COVID-19 disease susceptibility and severity which deserves further study. Three review papers on the topic have concluded there is either no difference or a reduced risk when looking at mortality and/or severe disease<sup>8,37,38</sup>. Our study builds on these previous reports where other standard statistical models were used for analysis, with potential introduction of biases<sup>11</sup>. Integrating standard cox models into a multi-state framework allows the study of separate outcomes simultaneously and allows the calculation of the transition probabilities, adding a layer of interpretation. We used a time-to-event analysis considering competing risks to account appropriately for censoring<sup>39</sup>, thereby robustly showing a modest, yet significant, positive effect of ACEI/ARBs use in recovery and survival of hospitalized COVID-19 patients accounting for confounding factors.

Our study uses nationwide hospital surveillance data, with mandatory participation, and includes all general hospitals (including university hospitals) in Belgium, both those managed by a public authority and privately managed are represented. The surveillance does not cover psychiatric hospitals or specialist hospitals<sup>14</sup>. The use of comprehensive datasets, but more so the utilization of models adequately fitting to time-to-event hospital data with mutually exclusive health states results in less probability of introducing biases and are crucial for correct evidence-based information for

1  
2  
3 381 decision making. Our study makes some assumptions, and unknowns such as the lack of information  
4  
5 382 on ACEI/ARBs exact indication and whether their use was continued after admission. Our models are  
6  
7 383 not adjusted for severity at baseline since we reasoned that hospital admission was already an  
8  
9 384 indicator of severe disease and 94% of patients had symptoms compatible with COVID-19 diagnosis at  
10  
11 385 baseline. Further, even though other events potentially indicating severity (ECMO, ARDS, pneumonia,  
12  
13 386 bacterial and/or fungal co-infection) were available in the database, only transfer to ICU was linked to  
14  
15 387 a calendar date and was therefore the only event which could be used as a proxy for severe health  
16  
17 388 state in our time-dependent model, indicating that our estimates might represent more a critical state  
18  
19 389 of the patient. In addition, admission to intensive care is not solely based on the clinical status of the  
20  
21 390 patient, but also on other criteria such as frailty. Also, ICU admission criteria might have been more  
22  
23 391 restrictive in the peak period of the epidemic whilst certain ICU were overloaded. Because the  
24  
25 392 surveillance data is limited to the most important variables, we cannot discard the possibility of some  
26  
27 393 degree of residual confounding in our results. An important limitation of our main analysis is the  
28  
29 394 impossibility of adjusting our models for smoking status, obesity and cognitive issues at baseline. Using  
30  
31 395 available smoking information was not considered due to the excessive number of missing values, and  
32  
33 396 the lack of information of the reason for the incomplete data. We used however data on obesity and  
34  
35 397 cognitive issues, which collection was introduced later, in a complete case analysis to confirm the  
36  
37 398 results obtained in the main model. Nevertheless, these analyses on a reduced sample of patients  
38  
39 399 should be interpreted with caution as a time effect is likely present because of the late data collection.  
40  
41 400 Finally, our analyses are based on patient's medical files and rely on how clinicians reported clinical  
42  
43 401 observations and anamnesis which might vary across hospitals, and are representative of the first so  
44  
45 402 called wave of the epidemic in Belgium, and associations might differ in subsequent studies.  
46  
47  
48  
49  
50  
51  
52  
53

54 404 **6. CONCLUSIONS**

55 405  
56  
57  
58  
59  
60

After adjustment for important confounders there is modest, yet significant, positive effect of ACEI/ARBs use on recovery and survival of hospitalized COVID-19 patients, without affecting admission to intensive care. This supports the use of ACEI/ARB in those patients who need them, also when needing hospitalization from COVID-19. These findings are based on an analytical model that adequately fits hospital data, where patients progress across different, competing, health states providing a more complete and accurate view of the research question within a reduced risk of bias framework. Integrating standard cox models into a multi-state framework allows the study of separate outcomes simultaneously and allows the calculation of the transition probabilities, adding a layer of interpretation. Multi-state models should be favoured over separate survival analysis when competing risks are present, and traditional methods such as logit functions should be discouraged when time-to-event is available.

#### **Contributorship statement**

JLP, MvdS, MAW, DV conceptualized the study. DV performed data curation and provided data. JLP, EG, EM, DS developed methodology, and performed analysis. JLP supervised the study, and drafted the manuscript. All authors have critically reviewed, commented, and approved the manuscript before submission.

#### **Competing interest**

The authors declare that they have no competing interests.

#### **Funding statement**

This research received no specific grant from any funding agency in the public, commercial or not-for-profit sectors.

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

**Data availability**

The data that support the findings of this study are available from Sciensano but restrictions apply to the availability of these data, which were used under license for the current study, and so are not publicly available. Data are however available from the authors upon reasonable request and with permission of Sciensano.

**Acknowledgments**

The authors thank the Members of the Belgian Collaborative Group on COVID-19 Hospital Surveillance: Samy Amir Aouachria, Centre Hospitalier Chrétien, Liège, Belgium; Kristof Bafort, Mariaziekenhuis, Pelt, Belgium; Leïla Belkhir, Cliniques Universitaires Saint-Luc, Brussels, Belgium; Koen Blot, Sciensano, Belgium; Nathalie Bossuyt, Sciensano, Belgium; Vincent Colombie, Centre Hospitalier Epicura, Baudour, Belgium; Nicolas Dauby, Centre Hospitalier Universitaire Saint-Pierre, Brussels, Belgium; Paul De Munter , Universitair Ziekenhuis Leuven, Leuven, Belgium; Robby De Pauw, Sciensano, Brussels, Belgium; Didier Delmarcelle, Clinique St. Jean, Brussels, Belgium; Mélanie Delvallee, Centre Hospitalier de Wallonie Picarde, Tournai, Belgium; Rémy Demeester, Centre Hospitalier Universitaire de Charleroi, Charleroi, Belgium; Thierry Dugernier, Clinique Saint-Pierre, Ottignies, Belgium; Xavier Holemans, Grand Hôpital de Charleroi, Charleroi, Belgium; Benjamin Kerzmann, Clinique Notre Dame de Grâce, Gosselies, Belgium; Pierre Yves Machurot, Centre Hospitalier de l'Ardenne, Belgium; Philippe Minette, Centres Hospitaliers Jolimont, Belgium; Jean-Marc Minon, Centre Hospitalier Régional de la Citadelle, Liège, Belgium; Saphia Mokrane, Hôpitaux Iris Sud, Brussels, Belgium; Catherine Nachtergal, Cliniques de l'Europe, Brussels, Belgium; Séverine Noirhomme, Centre Hospitalier Régional de Namur, Belgium; Denis Piérard, Universitair Ziekenhuis Brussel, Brussels; Camelia Rossi, Centre Hospitalier Universitaire Ambroise Paré, Mons, Belgium; Carole Schirvel, CHIREC, Brussels, Belgium; Erica Sermijn, A.S.Z. Ziekenhuis, Aalst, Belgium; Ben Serrien, Sciensano, Belgium; Fabio Taccone, Hôpital Erasme, Brussels, Belgium; Frank Staelens, OLV Ziekenhuis, Aalst, Belgium; Filip Triest, Algemeen Ziekenhuis Sint Lucas, Gent, Belgium; Nina Van

458 Goethem, Sciensano, Brussels, Belgium; Jens Van Praet, Algemeen Ziekenhuis Sint Jan, Brugge-  
459 Oostende, Belgium; Anke Vanhoenacker, Ziekenhuisnetwerk Antwerpen, Belgium; Roeland  
460 Verstraete, Algemeen Ziekenhuis, Monica, Belgium; Elise Willems, Algemeen Ziekenhuis Nikolaas,  
461 Sint-Niklaas, Belgium; Chloé Wyndham-Thomas, Sciensano, Belgium.

462

For peer review only

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

463

For peer review only

## REFERENCES

- 1 Yang, X. *et al.* Clinical course and outcomes of critically ill patients with SARS-CoV-2 pneumonia in Wuhan, China: a single-centered, retrospective, observational study. *Lancet Respir Med* **8**, P485-P481, doi:10.1016/S2213-2600(20)30079-5 (2020).
- 2 Guan, W. J. *et al.* Clinical Characteristics of Coronavirus Disease 2019 in China. *N Engl J Med* **382**, 1708-1730, doi:10.1056/NEJMoa2002032 (2020).
- 3 Zhang, J. J. *et al.* Clinical characteristics of 140 patients infected with SARS-CoV-2 in Wuhan, China. *Allergy* **75**, 1730-1741, doi:10.1111/all.14238 (2020).
- 4 Kassir, R. Risk of COVID-19 for patients with obesity. *Obesity Reviews* **21**, e13034, doi:10.1111/obr.13034.
- 5 Petrilli, C. M. *et al.* Factors associated with hospital admission and critical illness among 5279 people with coronavirus disease 2019 in New York City: prospective cohort study. *BMJ* **360**, m1966, doi:10.1101/2020.04.08.20057794 (2020).
- 6 Wan, Y., Shang, J., Graham, R., Baric, R. S. & Li, F. Receptor Recognition by the Novel Coronavirus from Wuhan: an Analysis Based on Decade-Long Structural Studies of SARS Coronavirus. *Journal of Virology* **94**, e00127-00120, doi:10.1128/jvi.00127-20 (2020).
- 7 Fang, L., Karakiulakis, G. & Roth, M. Are patients with hypertension and diabetes mellitus at increased risk for COVID-19 infection? *Lancet Respir Med* **8**, e21, doi:10.1016/S2213-2600(20)30116-8 (2020).
- 8 Zhang, J., Wang, M., Ding, W. & Wan, J. The interaction of RAAS inhibitors with COVID-19: Current progress, perspective and future. *Life Sci* **257**, 118142, doi:10.1016/j.lfs.2020.118142 (2020).
- 9 Mackey, K. *et al.* Risks and Impact of Angiotensin-Converting Enzyme Inhibitors or Angiotensin-Receptor Blockers on SARS-CoV-2 Infection in Adults. *Annals of Internal Medicine* **173**, 195-203, doi:10.7326/m20-1515 %m 32422062 (2020).
- 10 de Abajo, F. J. Renin-angiotensin system inhibitors and COVID-19: overwhelming evidence against an association. *The Lancet Digital Health* **3**, E70-E71, doi:10.1016/S2589-7500(20)30294-6 (2021).
- 11 Wolkewitz, M. *et al.* Statistical Analysis of Clinical COVID-19 Data: A Concise Overview of Lessons Learned, Common Errors and How to Avoid Them. *Clin Epidemiol* **12**, 925-928, doi:10.2147/CLEP.S256735 (2020).
- 12 Hazard, D. *et al.* Joint analysis of duration of ventilation, length of intensive care, and mortality of COVID-19 patients: a multistate approach. *BMC Medical Research Methodology* **20**, 206, doi:10.1186/s12874-020-01082-z (2020).
- 13 von Cube, M. *et al.* Harmonizing Heterogeneous Endpoints in Coronavirus Disease 2019 Trials Without Loss of Information. *Critical Care Medicine* **49**, e11-e19, doi:10.1101/2020.03.31.20049007 (2020).
- 14 Van Goethem, N. *et al.* Rapid establishment of a national surveillance of COVID-19 hospitalizations in Belgium. *Arch Public Health* **78**, 121, doi:10.1186/s13690-020-00505-z (2020).
- 15 WHO. COVID-19 and NCDs, <<https://www.who.int/internal-publications-detail/covid-19-and-ncds>> (2020).
- 16 de Azambuja, E. *et al.* Impact of solid cancer on in-hospital mortality overall and among different subgroups of patients with COVID-19: a nationwide, population-based analysis. *ESMO Open* **5**, e000947, doi:10.1136/esmoopen-2020-000947 (2020).
- 17 Catteau, L. *et al.* Low-dose hydroxychloroquine therapy and mortality in hospitalised patients with COVID-19: a nationwide observational study of 8075 participants. *International Journal of Antimicrobial Agents* **56**, 106144, doi:<https://doi.org/10.1016/j.ijantimicag.2020.106144> (2020).



1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

Survival R package. <<https://cran.r-project.org/web/packages/survival/survival.pdf>> (de Wreede, L. C., Fiocco, M. & Putter, H. The mstate package for estimation and prediction in non- and semi-parametric multi-state and competing risks models. *Comput Methods Programs Biomed* **99**, 261-274, doi:10.1016/j.cmpb.2010.01.001 (2010).

Mstate R package. <<https://cran.r-project.org/web/packages/mstate/mstate.pdf>> (Taccone, F. S. *et al.* The role of organizational characteristics on the outcome of COVID-19 patients admitted to the ICU in Belgium. *The Lancet Regional Health - Europe* **0**, 100019, doi:<https://doi.org/10.1016/j.lanepe.2020.100019> (2020).

Peng, Y. D. *et al.* [Clinical characteristics and outcomes of 112 cardiovascular disease patients infected by 2019-nCoV]. *Zhonghua Xin Xue Guan Bing Za Zhi* **48**, 450-455, doi:10.3760/cma.j.cn112148-20200220-00105 (2020).

Huang, Z. *et al.* The effect of RAS blockers on the clinical characteristics of COVID-19 patients with hypertension. *Ann Transl Med* **8**, 430, doi:10.21037/atm.2020.03.229 (2020).

Li, J., Wang, X., Chen, J., Zhang, H. & Deng, A. Association of Renin-Angiotensin System Inhibitors With Severity or Risk of Death in Patients With Hypertension Hospitalized for Coronavirus Disease 2019 (COVID-19) Infection in Wuhan, China. *JAMA Cardiol* **5**, 825-830, doi:10.1001/jamacardio.2020.1624 (2020).

Meng, X. *et al.* Angiotensin converting enzyme inhibitors and angiotensin receptor blockers improved the outcome of patients with severe COVID-19 and hypertension. *Sci China Life Sci* **1-4**, doi:10.1007/s11427-020-1813-0 (2020).

Feng, Y. *et al.* COVID-19 with Different Severities: A Multicenter Study of Clinical Features. *Am J Respir Crit Care Med* **201**, 1380-1388, doi:10.1164/rccm.202002-0445OC (2020).

Yang, G. *et al.* Effects of Angiotensin II Receptor Blockers and ACE (Angiotensin-Converting Enzyme) Inhibitors on Virus Infection, Inflammatory Status, and Clinical Outcomes in Patients With COVID-19 and Hypertension: A Single-Center Retrospective Study. *Hypertension* **76**, 51-58, doi:10.1161/HYPERTENSIONAHA.120.15143 (2020).

Mancia, G., Rea, F., Ludergnani, M., Apolone, G. & Corrao, G. Renin-Angiotensin-Aldosterone System Blockers and the Risk of Covid-19. *N Engl J Med* **382**, 2431-2440, doi:10.1056/NEJMoa2006923 (2020).

de Abajo, F. J. *et al.* Use of renin-angiotensin-aldosterone system inhibitors and risk of COVID-19 requiring admission to hospital: a case-population study. *Lancet* **395**, 1705-1714, doi:10.1016/S0140-6736(20)31030-8 (2020).

Reynolds, H. R. *et al.* Renin-Angiotensin-Aldosterone System Inhibitors and Risk of Covid-19. *N Engl J Med* **382**, 2441-2448, doi:10.1056/NEJMoa2008975 (2020).

Jung, S. Y., Choi, J. C., You, S. H. & Kim, W. Y. Association of renin-angiotensin-aldosterone system inhibitors with COVID-19-related outcomes in Korea: a nationwide population-based cohort study. *Clin Infect Dis* **71**, 2121-2128, doi:10.1093/cid/ciaa624 (2020).

Trifiro, G. *et al.* Renin-Angiotensin-Aldosterone System Inhibitors and Risk of Death in Patients Hospitalised with COVID-19: A Retrospective Italian Cohort Study of 43,000 Patients. *Drug Saf* **1-12**, doi:10.1007/s40264-020-00994-5 (2020).

Zhang, P. *et al.* Association of Inpatient Use of Angiotensin-Converting Enzyme Inhibitors and Angiotensin II Receptor Blockers With Mortality Among Patients With Hypertension Hospitalized With COVID-19. *Circulation Research* **126**, 1671-1681, doi:10.1161/CIRCRESAHA.120.317134 (2020).

Gao, C. *et al.* Association of hypertension and antihypertensive treatment with COVID-19 mortality: a retrospective observational study. *Eur Heart J* **41**, 2058-2066, doi:10.1093/eurheartj/ehaa433 (2020).

Fosbol, E. L. *et al.* Association of Angiotensin-Converting Enzyme Inhibitor or Angiotensin Receptor Blocker Use With COVID-19 Diagnosis and Mortality. *JAMA* **324**, 168-177, doi:10.1001/jama.2020.11301 (2020).

- 564 36 Hippisley-Cox, J. *et al.* Risk of severe COVID-19 disease with ACE inhibitors and angiotensin  
565 receptor blockers: cohort study including 8.3 million people. *Heart* **106**, 1503-1511,  
566 doi:10.1136/heartjnl-2020-317393 (2020).
- 567 37 Fernandez-Ruiz, I. RAAS inhibitors do not increase the risk of COVID-19. *Nat Rev Cardiol* **17**,  
568 383, doi:10.1038/s41569-020-0401-0 (2020).
- 569 38 Jarcho, J. A., Ingelfinger, J. R., Hamel, M. B., D'Agostino, R. B., Sr. & Harrington, D. P. Inhibitors  
570 of the Renin-Angiotensin-Aldosterone System and Covid-19. *N Engl J Med* **382**, 2462-2464,  
571 doi:10.1056/NEJMe2012924 (2020).
- 572 39 Putter, H., Fiocco, M. & Geskus, R. B. Tutorial in biostatistics: competing risks and multi-state  
573 models. *Stat Med* **26**, 2389-2430, doi:10.1002/sim.2712 (2007).
- 574
- 575

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

**ACKNOWLEDGMENTS**

The authors thank the Members of the Belgian Collaborative Group on COVID-19 Hospital Surveillance: Samy Amir Aouachria, Centre Hospitalier Chrétien, Liège, Belgium; Kristof Bafort, Mariaziekenhuis, Pelt, Belgium; Leïla Belkhir, Cliniques Universitaires Saint-Luc, Brussels, Belgium; Koen Blot, Sciensano, Belgium; Nathalie Bossuyt, Sciensano, Belgium; Vincent Colombie, Centre Hospitalier Epicura, Baudour, Belgium; Nicolas Dauby, Centre Hospitalier Universitaire Saint-Pierre, Brussels, Belgium; Paul De Munter, Universitair Ziekenhuis Leuven, Leuven, Belgium; Robby De Pauw, Sciensano, Brussels, Belgium; Didier Delmarcelle, Clinique St. Jean, Brussels, Belgium; Mélanie Delvallee, Centre Hospitalier de Wallonie Picarde, Tournai, Belgium; Rémy Demeester, Centre Hospitalier Universitaire de Charleroi, Charleroi, Belgium; Thierry Dugernier, Clinique Saint-Pierre, Ottignies, Belgium; Xavier Holemans, Grand Hôpital de Charleroi, Charleroi, Belgium; Benjamin Kerzmann, Clinique Notre Dame de Grâce, Gosselies, Belgium; Pierre Yves Machurot, Centre Hospitalier de l'Ardenne, Belgium; Philippe Minette, Centres Hospitaliers Jolimont, Belgium; Jean-Marc Minon, Centre Hospitalier Régional de la Citadelle, Liège, Belgium; Saphia Mokrane, Hôpitaux Iris Sud, Brussels, Belgium; Catherine Nachtergal, Cliniques de l'Europe, Brussels, Belgium; Séverine Noirhomme, Centre Hospitalier Régional de Namur, Belgium; Denis Piérard, Universitair Ziekenhuis Brussel, Brussels; Camelia Rossi, Centre Hospitalier Universitaire Ambroise Paré, Mons, Belgium; Carole Schirvel, CHIREC, Brussels, Belgium; Erica Sermijn, A.S.Z. Ziekenhuis, Aalst, Belgium; Ben Serrien, Sciensano, Belgium; Fabio Taccone, Hôpital Erasme, Brussels, Belgium; Frank Staelens, OLV Ziekenhuis, Aalst, Belgium; Filip Triest, Algemeen Ziekenhuis Sint Lucas, Gent, Belgium; Nina Van Goethem, Sciensano, Brussels, Belgium; Jens Van Praet, Algemeen Ziekenhuis Sint Jan, Brugge-Oostende, Belgium; Anke Vanhoenacker, Ziekenhuisnetwerk Antwerpen, Belgium; Roeland Verstraete, Algemeen Ziekenhuis, Monica, Belgium; Elise Willems, Algemeen Ziekenhuis Nikolaas, Sint-Niklaas, Belgium; Chloé Wyndham-Thomas, Sciensano, Belgium.

## **AUTHOR CONTRIBUTIONS**

JLP, MvdS, MAW, DV conceptualized the study. DV performed data curation and provided data. JLP, EG, EM, DS developed methodology, and performed analysis. JLP supervised the study, and drafted the manuscript. All authors have critically reviewed, commented, and approved the manuscript before submission.

## **DATA AVAILABILITY**

The data that support the findings of this study are available from Sciensano but restrictions apply to the availability of these data, which were used under license for the current study, and so are not publicly available. Data are however available from the authors upon reasonable request and with permission of Sciensano.

## **ETHICS DECLARATION**

Ethical clearance was obtained from the Institutional Review Board (IRB) of the Institute of Tropical Medicine after revision of the research protocol (num. 1393/20, 02/05/2020).

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

**FIGURE TITLES**

**Figure 1.** Schematic representation of the competing risk multi-state model and transition event matrix (number (%) patients in each transition.

**Figure 2.** Plots for cumulative transition hazards (A), and state transition probabilities (B), and transition probabilities after transfer to intensive care (C) in a multi-state competing risk model considering ACEI/ARBs use (dashed line) versus no use (solid line).

**Figure 1 Supplemental.** Staked probability plot of the state-occupation probabilities in a multi-state competing risk model considering ACEI/ARBs use (A) versus no use (B).

**Table 1.** COVID-19 patient's characteristics at hospital admission according to ACEI/ARBs use

	Total	ACEI/ARBs		
	(n = 11717)	No use (n = 8189, 69.9%)	Use (n = 2810, 23.9%)	Unknown use (n = 718, 6.1%)
<b>Demographics</b>				
Age (years) (mean (SD))	67.82 (17.17)	65.70 (17.90)	74.08 (12.85)	67.47 (17.50)
Equal or more than 70 years old (n, %)	6044 (51.6)	3791 (46.3)	1886 (67.1)	367 (51.1)
Sex (n, % males)	6154 (52.5)	4227 (51.6)	1562 (55.6)	365 (50.8)
Missing (n, %)	129 (1.1)	93 (1.1)	25 (0.9)	11 (1.5)
<b>Risk factors</b>				
Smokers (n, %)	606 (5.2)	440 (5.4)	142 (5.1)	24 (3.3)
Missing (n, %)	5413 (46.2)	3667 (44.8)	1160 (41.3)	586 (81.6)
Flu vaccination (n, %)	841 (7.2)	572 (7.0)	250 (8.9)	19 (2.6)
Missing (n, %)	10076 (86.0)	7018 (85.7)	2374 (84.5)	684 (95.3)
Obesity (n, %)*	782 (6.7)	478 (5.8)	271 (9.6)	33 (4.6)
Missing (n, %)	3887 (33.2)	2735 (33.4)	870 (31.0)	282 (39.3)
<b>Chronic comorbidities</b>				
HBP (n, %)	4593 (39.2)	2343 (28.6)	2090 (74.4)	160 (22.3)
DM (n, %)	2522 (21.5)	1486 (18.1)	936 (33.3)	100 (13.9)
Chronic renal disease (n, %)	1513 (12.9)	911 (11.1)	541 (19.3)	61 (8.5)
CVD (n, %)	3984 (34.0)	2326 (28.4)	1493 (53.1)	165 (23.0)
Chronic lung disease (n, %)	1731 (14.8)	1180 (14.4)	473 (16.8)	78 (10.9)
Cognitive impairment (n, %)**	1320 (11.3)	922 (11.3)	331 (11.8)	67 (9.3)
Missing (n, %)	668 (5.7)	461 (5.6)	173 (6.2)	34 (4.7)
Chronic neuro-muscular disease (n, %)	993 (8.5)	704 (8.6)	241 (8.6)	48 (6.7)
Solid malignant neoplasms (n, %)	990 (8.4)	697 (8.5)	261 (9.3)	32 (4.5)
Chronic liver disease (n, %)	301 (2.6)	210 (2.6)	79 (2.8)	12 (1.7)
Immunodepression (n, %)	297 (2.5)	224 (2.7)	64 (2.3)	9 (1.3)
Hematological cancers (n, %)	216 (1.8)	154 (1.9)	56 (2.0)	6 (0.8)
<b>Combination of comorbidities</b>				
None (n, %)	4760 (40.6)	4145 (50.6)	192 (6.8)	423 (58.9)
CVD & HBP (n, %)	1386 (11.8)	713 (8.7)	633 (22.5)	41 (5.7)
CVD & DM (n, %)	385 (3.3)	248 (3.0)	113 (4.0)	24 (3.3)
HBP & DM (n, %)	682 (5.8)	348 (4.2)	309 (11.0)	25 (3.5)
CVD & HBP & DM (n, %)	401 (50.6)	401 (4.9)	423 (15.1)	20 (2.8)

ACEI/ARBs, Angiotensin converting enzyme inhibitors / Angiotensin receptor blockers; CVD, cardiovascular disease; DM, diabetes mellitus; HBP, high blood pressure. \*Values only collected after April 3, 2020. \*\*Values only reported after March 23, 2020.

**Table 2.** Frequency of severity events among COVID-19 patients during hospital stay and recorded outcomes at discharge according to ACEI/ARBs use at admission

		ACEI/ARBs			
		Total	No use	Use	Unknown use
		(n = 11717)	(n = 8189, 69.9%)	(n = 2810, 23.9%)	(n = 718, 6.1%)
<b>Severe conditions</b>					
Pneumonia (n, %)		9265 (79.1)	6501 (79.4)	2260 (80.4)	504 (70.2)
Missing (n, %)		532 (4.5)	372 (4.5)	73 (73)	87 (87)
Superinfection (n, %)		2268 (19.4)	1548 (18.9)	589 (21.0)	131 (18.2)
Missing (n, %)		1277 (10.9)	820 (10.0)	320 (11.4)	137 (19.1)
ARDS (n, %)		1492 (12.7)	996 (12.2)	389 (13.8)	107 (14.9)
Missing (n, %)		1047 (8.9)	671 (8.2)	263 (9.4)	113 (15.7)
Mechanical ventilation (n, %)		893 (7.6)	571 (7.0)	249 (8.9)	73 (10.2)
Missing (n, %)		627 (5.4)	383 (4.7)	170 (6.0)	74 (10.3)
Number of severe conditions (n, %)					
	None	2143 (18.3)	1492 (18.2)	465 (16.5)	186 (25.9)
	One	6537 (55.8)	4620 (56.4)	1565 (55.7)	352 (49.0)
	Two or more	3037 (25.9)	2077 (25.4)	780 (27.8)	180 (25.1)
<b>Intensive care</b>					
Transfer to ICU (n, %)		1518 (13.0)	990 (12.1)	425 (15.1)	103 (14.3)
Missing (n, %)		16 (0.1)	16 (0.2)	0 (0)	0 (0)
Transfer to ICU + pneumonia (n, %)		1423 (93.7)	932 (94.1)	395 (92.9)	96 (93.2)
Transfer to ICU + superinfection (n, %)		653 (43.0)	423 (42.7)	181 (42.6)	49 (47.6)
Transfer to ICU + ARDs (n, %)		831 (54.7)	547 (55.3)	224 (52.7)	60 (58.3)
Transfer to ICU + mechanical ventilation (n, %)		880 (58.0)	561 (56.7)	246 (57.9)	73 (70.9)
Length (days) of ICU stay (mean (SD))		11.5 (10.7)	11.415 (10.7)	11.3 (10.8)	12.9 (11.0)
<b>Discharge status</b>					
Recovered at discharge (n, %)		6003 (51.2)	4244 (51.8)	1378 (49.0)	381 (53.1)
Recovered at home (n, %)		3093 (26.4)	2201 (26.9)	722 (25.7)	170 (23.7)
In-hospital death (n, %)		2388 (20.4)	1574 (19.2)	622 (23.6)	152 (21.2)
Transferred (n, %)		201 (1.7)	149 (51.8)	44 (49.0)	8 (53.1)
Unknown (n, %)		32 (0.3)	21 (0.3)	4 (0.1)	7 (1.0)
Length (days) of hospital stay (mean (SD))		12.6 (10.9)	12.1 (10.5)	13.9 (11.7)	12.2 (11.6)

ACEI/ARBs, Angiotensin converting enzyme inhibitors / Angiotensin receptor blockers; ARDS, acute respiratory distress syndrome; ICU, intensive care unit.


**Table 3.** State-arrival extended Cox-Markov multivariate model's transitions hazard ratios (HR, 95%CI) as a function of ACEI/ARBs

Transition		ACEI/ARBs use			
		Model 1	p-value	Model 2	p-value
1	Admission → Severity	1.15 (0.98, 1.36)	0.092	1.10 (0.88, 1.36)	0.409
2	Admission → Recovery	<b>1.07 (1.01, 1.13)</b>	0.027	1.05 (0.98, 1.13)	0.182
3	Admission → Death	<b>0.83 (0.75, 0.93)</b>	0.001	<b>0.80 (0.70, 0.91)</b>	0.001
4	Severity → Recovery	1.16 (0.97, 1.38)	0.098	1.16 (0.93, 1.45)	0.195
5	Severity → Death	0.91 (0.73, 1.13)	0.381	1.11 (0.83, 1.49)	0.485

ACEI/ARBs, Angiotensin converting enzyme inhibitors / Angiotensin receptor blockers Model 1: Adjusted for gender, age (years), prevalent CVD, HBP, DM, and time (days) to severity; Model 2: Further by prevalent obesity, and cognitive issues. Full output is presented in Table 3 supplemental.




Figure 1.



From	To	Admission	Severity	Recovery	Death	No event	Total
Admission		0	863 (8.4) <sup>1</sup>	7633 (73.9) <sup>2</sup>	1738 (16.8) <sup>3</sup>	93 (0.9)	10327
Severity		0	0	817 (58.3) <sup>4</sup>	468 (33.4) <sup>5</sup>	117 (8.3)	1402*
Recovery		0	0	0	0	8450 (100)	8450
Death		0	0	0	0	2206 (100)	2206

Numbers in superscript represent transitions depicted in the figure. \*539 patients were directly admitted to ICU

**Figure 2.**

SUPPLEMENTAL MATERIAL

**Table 1 Supplemental.** Frequency of symptoms and clinical signs compatible with COVID-19 at hospital admission according to ACEI/ARBs use

	Total	ACEI/ARBs		
		No use	Use	Unknown use
	(n = 11717)	(n = 8189, 69.9%)	(n = 2810, 23.9%)	(n = 718, 6.1%)
<b>Symptoms at admission</b>	11039 (94.2)	7705 (94.1)	2673 (95.1)	661 (92.1)
Fever (n, %)	7187 (61.3)	5112 (62.4)	1653 (58.8)	422 (58.8)
Cough (n, %)	6231 (53.2)	4444 (54.3)	1438 (51.2)	349 (48.6)
Breathlessness (n, %)	5997 (51.2)	4194 (51.2)	1349 (52.0)	376 (47.6)
Weakness (n, %)	4672 (39.9)	3253 (39.7)	1153 (41.0)	266 (37.0)
Pain (n, %)	2622 (22.4)	1931 (23.6)	544 (19.4)	147 (20.5)
Diarrhea (n, %)	1673 (14.3)	1173 (14.3)	438 (15.6)	62 (8.6)
Nausea and vomiting (n, %)	1364 (11.6)	987 (12.1)	311 (11.1)	66 (9.2)
Headache (n, %)	1234 (10.5)	932 (11.4)	245 (8.7)	57 (7.9)
Irritability (n, %)	838 (7.2)	556 (6.8)	243 (8.6)	39 (5.4)
Throat pain (n, %)	700 (6.0)	520 (6.3)	141 (5.0)	39 (5.4)
Anosmia (n, %)	424 (3.6)	334 (4.1)	79 (2.8)	11 (1.5)
Missing (n, %)	855 (7.3)	593 (7.2)	216 (7.7)	46 (6.4)
Runny nose (n, %)	416 (3.6)	301 (3.7)	91 (3.2)	24 (3.3)
<b>Clinical signs at admission</b>	9993 (85.3)	7005 (85.5)	2452 (87.3)	536 (74.7)
Abnormal pulmonary imaging (n, %)*	7396 (63.1)	5271 (64.4)	1835 (65.3)	290 (40.4)
Abnormal pulmonary auscultation (n, %)	5245 (44.8)	3701 (45.2)	1343 (47.8)	201 (28.0)
Dyspnea (n, %)	4966 (42.4)	3462 (42.3)	1196 (42.6)	308 (42.9)
Pharyngitis (n, %)	244 (2.1)	174 (2.1)	58 (2.1)	12 (1.7)
Coma (n, %)	72 (0.6)	50 (0.6)	16 (0.6)	6 (0.8)
Conjunctivitis (n, %)	57 (0.5)	37 (0.5)	18 (0.6)	2 (0.3)
Convulsions (n, %)	15 (0.1)	12 (0.1)	3 (0.1)	0 (0.0)

ACEI; Angiotensin converting enzyme inhibitors, ARBs. Angiotensin receptor blockers

\* Reported as abnormal pulmonary imaging compatible with pneumonia

**Table 2 Supplemental.** Results (OR, 95%CI) of variable selection models of conditions associated with ACEI/ARBs use among factors with COVID-19 prognosis<sup>1</sup>

Variable	Model 1	p-value	Model 2	p-value
Intercept	0.04 (0.03-0.05)	< 0.0001	0.03 (0.03-0.05)	< 0.001
Gender (male)	1.33 (1.21-1.47)	< 0.0001	1.26 (1.12-1.41)	< 0.001
Age	1.01 (1.01-1.02)	< 0.0001	1.01 (1.01-1.02)	< 0.001
CVD	1.71 (1.55-1.90)	< 0.0001	1.65 (1.46-1.87)	< 0.001
Diabetes	1.38 (1.24-1.54)	< 0.0001	1.37 (1.20-1.56)	< 0.001
HBP	5.65 (5.10-6.27)	< 0.0001	5.28 (4.66-6.00)	< 0.001
Obesity			1.33 (1.10-1.59)	< 0.01
Cognitive impairment			0.69 (0.58-0.82)	< 0.001

<sup>1</sup> Backwards stepwise logistic regression with variable selection according to AIC; OR, odds ratio. Model 1 included: Gender, age, HBP, CVD, DM, chronic renal disease, chronic liver disease, chronic lung disease, solid malignant neoplasms, hematological cancers, immunodepression (N=10866); Model 2 included: Model 1 variables plus cognitive impairment, chronic neuro-muscular disease, and obesity, for which a high degree of missingness was observed and represents a complete case analysis (N=7,294)

**Table 3 Supplemental.** State-arrival extended Cox-Markov models transitions hazard ratios (HR, 95%CI) as a function of ACEI/ARBs and identified confounders


**Model 1**

Transition		ACEI/ARBs use	Male gender	Age < 70	Age > 70	CVD	HBP	DM	Time to severity	
1	Admission to Severity	1.15 (0.98, 1.36)	<b>1.83 (1.58, 2.12)</b>	<b>1.02 (1.01, 1.02)</b>	<b>0.90 (0.89, 0.91)</b>	0.96 (0.82, 1.12)	<b>1.35 (1.15, 1.58)</b>	0.94 (0.79, 1.11)		
		0.092	<0.0001	<0.0001	<0.0001	0.575	<0.0001	0.440		
2	Admission to Recovery	<b>1.07 (1.01, 1.13)</b>	<b>0.89 (0.85, 0.93)</b>	<b>0.97 (0.97, 0.97)</b>	<b>0.96 (0.96, 0.97)</b>	<b>0.86 (0.81, 0.91)</b>	<b>0.95 (0.89, 1.00)</b>	<b>0.92 (0.87, 0.98)</b>		
		0.027	<0.0001	<0.0001	<0.0001	<0.0001	0.043	0.007		
3	Admission to Death	<b>0.83 (0.75, 0.93)</b>	<b>1.45 (1.31, 1.59)</b>	<b>1.11 (1.09, 1.13)</b>	<b>1.05 (1.04, 1.05)</b>	<b>1.13 (1.02, 1.25)</b>	1.00 (0.90, 1.10)	1.09 (0.98, 1.22)		
		0.001	<0.0001	<0.0001	<0.0001	0.015	0.947	0.102		
4	Severity to recovery	1.16 (0.97, 1.38)	0.90 (0.77, 1.04)	<b>0.97 (0.96, 0.97)</b>	1.00 (0.98, 1.02)	1.03 (0.87, 1.21)	0.95 (0.81, 1.12)	1.01 (0.85, 1.19)	<b>0.95 (0.93, 0.97)</b>	
		0.098	0.158	<0.0001	0.799	0.766	0.570	0.919	<0.0001	
5	Severity to death	0.91 (0.73, 1.13)	1.11 (0.91, 1.36)	<b>1.05 (1.03, 1.06)</b>	1.05 (1.03, 1.07)	1.07 (0.88, 1.30)	0.90 (0.74, 1.11)	<b>1.25 (1.02, 1.53)</b>	1.02 (1.00, 1.04)	
		0.381	0.316	<0.0001	<0.0001	0.494	0.326	0.034	0.056	

**Model 2**

Transition		ACEI/ARBs use	Male gender	Age < 70	Age > 70	CVD	HBP	DM	Cognitive issues	Obesity	Time to severity
1	Admission to severity	1.10 (0.88, 1.36)	<b>1.71 (1.42, 2.08)</b>	<b>1.02 (1.01, 1.03)</b>	<b>0.92 (0.90, 0.93)</b>	0.91 (0.74, 1.12)	<b>1.45 (1.18, 1.79)</b>	0.96 (0.77, 1.19)	<b>0.52 (0.35, 0.77)</b>	<b>1.58 (1.24, 2.02)</b>	
		0.409	<0.0001	0.001	<0.0001	0.361	0.001	0.686	0.001	<0.0001	
2	Admission to recovery	1.05 (0.98, 1.13)	<b>0.89 (0.84, 0.94)</b>	<b>0.97 (0.97, 0.97)</b>	<b>0.97 (0.96, 0.97)</b>	<b>0.88 (0.82, 0.94)</b>	0.97 (0.91, 1.04)	0.93 (0.87, 1.00)	<b>0.83 (0.76, 0.91)</b>	1.01 (0.92, 1.11)	
		0.182	<0.0001	<0.0001	<0.0001	<0.0001	0.377	0.061	<0.0001	0.882	
3	Admission to death	<b>0.80 (0.70, 0.91)</b>	<b>1.48 (1.31, 1.66)</b>	<b>1.10 (1.07, 1.12)</b>	<b>1.05 (1.04, 1.05)</b>	<b>1.13 (1.00, 1.27)</b>	1.03 (0.91, 1.16)	1.11 (0.97, 1.27)	<b>1.36 (1.19, 1.55)</b>	0.97 (0.76, 1.24)	
		0.001	<0.0001	<0.0001	<0.0001	0.048	0.685	0.116	<0.0001	0.824	
4	Severity to recovery	1.16 (0.93, 1.45)	0.90 (0.75, 1.09)	<b>0.97 (0.96, 0.98)</b>	1.00 (0.97, 1.02)	1.12 (0.91, 1.38)	1.00 (0.81, 1.24)	0.95 (0.77, 1.16)	1.22 (0.84, 1.77)	0.97 (0.77, 1.22)	<b>0.94 (0.91, 0.97)</b>
		0.195	0.272	<0.0001	0.861	0.288	0.984	0.597	0.297	0.781	<0.0001
5	Severity to death	1.11 (0.83, 1.49)	1.22 (0.93, 1.60)	<b>1.04 (1.02, 1.06)</b>	<b>1.06 (1.03, 1.08)</b>	1.08 (0.82, 1.41)	0.81 (0.61, 1.08)	<b>1.44 (1.10, 1.90)</b>	0.89 (0.53, 1.50)	0.97 (0.69, 1.36)	<b>1.04 (1.01, 1.06)</b>
		0.485	0.146	<0.0001	<0.0001	0.591	0.150	0.008	0.668	0.860	0.003

ACEI/ARBs, Angiotensin converting enzyme inhibitors / Angiotensin receptor blockers; CVD, cardiovascular disease; DM, diabetes mellitus; HBP, high blood pressure. Model 1: Adjusted for gender, age (years), CVD, HBP, DM, and time (days) to severity; Model 2: Further by prevalent obesity, and cognitive issues.

**Figure 1 Supplemental.**

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

STROBE Statement—checklist of items that should be included in reports of observational studies

	Item No	Recommendation	Page No
Title and abstract	1	(a) Indicate the study’s design with a commonly used term in the title or the abstract	1
		(b) Provide in the abstract an informative and balanced summary of what was done and what was found	2
Introduction			
Background/rationale	2	Explain the scientific background and rationale for the investigation being reported	5
Objectives	3	State specific objectives, including any prespecified hypotheses	6
Methods			
Study design	4	Present key elements of study design early in the paper	7
Setting	5	Describe the setting, locations, and relevant dates, including periods of recruitment, exposure, follow-up, and data collection	7,8
Participants	6	(a) Cohort study—Give the eligibility criteria, and the sources and methods of selection of participants. Describe methods of follow-up Case-control study—Give the eligibility criteria, and the sources and methods of case ascertainment and control selection. Give the rationale for the choice of cases and controls Cross-sectional study—Give the eligibility criteria, and the sources and methods of selection of participants	8,9
		(b) Cohort study—For matched studies, give matching criteria and number of exposed and unexposed Case-control study—For matched studies, give matching criteria and the number of controls per case	
Variables	7	Clearly define all outcomes, exposures, predictors, potential confounders, and effect modifiers. Give diagnostic criteria, if applicable	9
Data sources/ measurement	8*	For each variable of interest, give sources of data and details of methods of assessment (measurement). Describe comparability of assessment methods if there is more than one group	8,9
Bias	9	Describe any efforts to address potential sources of bias	9,10
Study size	10	Explain how the study size was arrived at	9
Quantitative variables	11	Explain how quantitative variables were handled in the analyses. If applicable, describe which groupings were chosen and why	10
Statistical methods	12	(a) Describe all statistical methods, including those used to control for confounding	9-11
		(b) Describe any methods used to examine subgroups and interactions	10
		(c) Explain how missing data were addressed	10
		(d) Cohort study—If applicable, explain how loss to follow-up was addressed Case-control study—If applicable, explain how matching of cases and controls was addressed Cross-sectional study—If applicable, describe analytical methods taking account of sampling strategy	10
		(e) Describe any sensitivity analyses	10

Continued on next page

**Results**

Participants	13*	(a) Report numbers of individuals at each stage of study—eg numbers potentially eligible, examined for eligibility, confirmed eligible, included in the study, completing follow-up, and analysed	10, and Fig 1
		(b) Give reasons for non-participation at each stage	10
		(c) Consider use of a flow diagram	Fig 1
Descriptive data	14*	(a) Give characteristics of study participants (eg demographic, clinical, social) and information on exposures and potential confounders	11-14, Table 1
		(b) Indicate number of participants with missing data for each variable of interest	25, Table 1
		(c) <i>Cohort study</i> —Summarise follow-up time (eg, average and total amount)	26, Table 2
Outcome data	15*	<i>Cohort study</i> —Report numbers of outcome events or summary measures over time	26, table 2
		<i>Case-control study</i> —Report numbers in each exposure category, or summary measures of exposure	
		<i>Cross-sectional study</i> —Report numbers of outcome events or summary measures	
Main results	16	(a) Give unadjusted estimates and, if applicable, confounder-adjusted estimates and their precision (eg, 95% confidence interval). Make clear which confounders were adjusted for and why they were included	27, table 3
		(b) Report category boundaries when continuous variables were categorized	All tables
		(c) If relevant, consider translating estimates of relative risk into absolute risk for a meaningful time period	
Other analyses	17	Report other analyses done—eg analyses of subgroups and interactions, and sensitivity analyses	27, Table 3

**Discussion**

Key results	18	Summarise key results with reference to study objectives	14-16
Limitations	19	Discuss limitations of the study, taking into account sources of potential bias or imprecision. Discuss both direction and magnitude of any potential bias	16-17
Interpretation	20	Give a cautious overall interpretation of results considering objectives, limitations, multiplicity of analyses, results from similar studies, and other relevant evidence	14-16
Generalisability	21	Discuss the generalisability (external validity) of the study results	14-16

**Other information**

Funding	22	Give the source of funding and the role of the funders for the present study and, if applicable, for the original study on which the present article is based	3
---------	----	---	---

\*Give information separately for cases and controls in case-control studies and, if applicable, for exposed and unexposed groups in cohort and cross-sectional studies.



1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

**Note:** An Explanation and Elaboration article discusses each checklist item and gives methodological background and published examples of transparent reporting. The STROBE checklist is best used in conjunction with this article (freely available on the Web sites of PLoS Medicine at <http://www.plosmedicine.org/>, Annals of Internal Medicine at <http://www.annals.org/>, and Epidemiology at <http://www.epidem.com/>). Information on the STROBE Initiative is available at [www.strobe-statement.org](http://www.strobe-statement.org).

For peer review only

# BMJ Open

## Insights into the association of ACEI/ARBs use and COVID-19 prognosis: A multi-state modelling study of nationwide hospital surveillance data from Belgium

Journal:	<i>BMJ Open</i>
Manuscript ID	bmjopen-2021-053393.R1
Article Type:	Original research
Date Submitted by the Author:	23-Aug-2021
Complete List of Authors:	Penalvo, Jose; Institute of Tropical Medicine Genbrugge, Els; Institute of Tropical Medicine Mertens, Elly; Institute of Tropical Medicine, Sagastume, Diana; Institute of Tropical Medicine van der Sande, Marianne A. B.; Institute of Tropical Medicine Widdowson, Marc-Alain; Institute of Tropical Medicine Van Beckhoven, Dominique; Sciensano
<b>Primary Subject Heading</b>:	Pharmacology and therapeutics
Secondary Subject Heading:	Epidemiology
Keywords:	COVID-19, Hypertension < CARDIOLOGY, STATISTICS & RESEARCH METHODS

SCHOLARONE™  
Manuscripts



I, the Submitting Author has the right to grant and does grant on behalf of all authors of the Work (as defined in the below author licence), an exclusive licence and/or a non-exclusive licence for contributions from authors who are: i) UK Crown employees; ii) where BMJ has agreed a CC-BY licence shall apply, and/or iii) in accordance with the terms applicable for US Federal Government officers or employees acting as part of their official duties; on a worldwide, perpetual, irrevocable, royalty-free basis to BMJ Publishing Group Ltd ("BMJ") its licensees and where the relevant Journal is co-owned by BMJ to the co-owners of the Journal, to publish the Work in this journal and any other BMJ products and to exploit all rights, as set out in our [licence](#).

The Submitting Author accepts and understands that any supply made under these terms is made by BMJ to the Submitting Author unless you are acting as an employee on behalf of your employer or a postgraduate student of an affiliated institution which is paying any applicable article publishing charge ("APC") for Open Access articles. Where the Submitting Author wishes to make the Work available on an Open Access basis (and intends to pay the relevant APC), the terms of reuse of such Open Access shall be governed by a Creative Commons licence – details of these licences and which [Creative Commons](#) licence will apply to this Work are set out in our licence referred to above.

Other than as permitted in any relevant BMJ Author's Self Archiving Policies, I confirm this Work has not been accepted for publication elsewhere, is not being considered for publication elsewhere and does not duplicate material already published. I confirm all authors consent to publication of this Work and authorise the granting of this licence.

**Insights into the association of ACEI/ARBs use and COVID-19 prognosis: A multi-state modelling study of nationwide hospital surveillance data from Belgium**

José L. Peñalvo<sup>1</sup>, Els Genbrugge<sup>1</sup>, Elly Mertens<sup>1</sup>, Diana Sagastume<sup>1</sup>, Marianne A. B. van der Sande<sup>2,3</sup>, Marc-Alain Widdowson<sup>4</sup>, Dominique Van Beckhoven<sup>5</sup>

<sup>1</sup> Non-Communicable Diseases Unit, Department of Public Health, Institute of Tropical Medicine, Antwerp, Belgium

<sup>2</sup> Department of Public Health, Institute of Tropical Medicine, Antwerp, Belgium

<sup>3</sup> Global Health Julius Centre for Health Sciences and Primary Care, University Medical Centre Utrecht, Utrecht, The Netherlands.

<sup>4</sup> Institute of Tropical Medicine, Antwerp, Belgium

<sup>5</sup> Department of Epidemiology and Public Health, Sciensano, Brussels, Belgium

**Correspondence**

José L. Peñalvo, Non-Communicable Diseases Unit, Department of Public Health, Institute of Tropical Medicine. Nationalestraat 150, 2000-Antwerp, Belgium. Tel: +32(0)32476251, email [jpenalvo@itg.be](mailto:jpenalvo@itg.be)

**Keywords**

Angiotensin-converting enzyme inhibitors (ACEI); Angiotensin-receptor blockers (ARBs); COVID-19; comorbidities; multi-state model.

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

**Abstract**

*Objectives:* The widespread use of angiotensin-converting enzyme inhibitors (ACEI) and angiotensin receptor blockers (ARB) by chronic patients raised early concerns on the potential exacerbation of COVID-19 severity and fatality. Previous studies addressing this question have used standard methods that may lead to biased estimates when analyzing hospital data because of the presence of competing events and event-related dependency. We investigated the association of ACEI/ARBs use with COVID-19 disease outcomes using time-to-event data in a multi-state setting to account for competing events and minimize bias.

*Setting:* Nationwide surveillance data from 119 Belgian hospitals.

*Participants:* Medical records of 10,866 patients hospitalised from March 14 to June 14, 2020 with a confirmed SARS-CoV-19 infection and information about ACEI/ARBs use.

*Primary outcome measure:* Multi-state, multivariate Cox-Markov models were used to estimate the hazards of patients transitioning through health states from admission to discharge or death, along with transition probabilities calculated by combining the baseline cumulative hazard and regression coefficients.

*Results:* After accounting for potential confounders there was no discernable association between ACEI/ARBs use and transfer to intensive care (ICU). Contrastingly, for patients without ICU transfer, ACEI/ARBs use was associated with a modest increase in recovery (HR 1.07 [95%CI 1.01-1.13],  $p=0.027$ ) and reduction in fatality (0.83, 0.75-0.93,  $p=0.001$ ) transitions. For patients transferred to ICU admission, no evidence of an association between ACEI/ARBs use and recovery (1.16, 0.97-1.38,  $p=0.098$ ) or in-hospital death (0.91, 0.73-1.12,  $p=0.381$ ) was observed. Male gender and older age were significantly associated with higher risk of ICU admission or death. Chronic cardiometabolic comorbidities were also associated with less recovery.

*Conclusions:* For the first time, a multistate model was used to address magnitude and direction of the association of ACEI/ARBs use on COVID-19 progression. By minimizing bias, this study provided a robust indication of a protective, albeit modest, association with recovery and survival.

**Keywords:** Angiotensin-converting enzyme inhibitors (ACEI); Angiotensin-receptor blockers (ARBs); COVID-19; comorbidities; multi-state model.

### **Strengths and limitations of this study**

- The study uses nationwide hospital surveillance data, and includes all general hospitals in Belgium.
- The use of a comprehensive database, but more so the utilization of models adequately fitting time-to-event data with mutually exclusive health states results in less probability of introducing biases and are crucial for correct evidence-based information for decision making.
- Only transfer to intensive care was linked to a calendar date and was therefore the only event which could be used as a proxy for severe disease state in our time-dependent model, indicating that our estimates might represent more a critical state of the patient.
- Information of ACEI/ARBs use was available at admission only, without any further information on the in-hospital use of ACEI/ARBs for those patients which could introduce a risk of immeasurable time bias if treatment discontinuation vs continuation has an impact on COVID-19 severity outcomes.

1. BACKGROUND

COVID-19 is known to affect more severely to older individuals, men and patients with chronic respiratory or cardiometabolic conditions such as cardiovascular disease (CVD), hypertension (HTN) and diabetes (DM) <sup>1-3</sup>. Also, common risk factors for chronic conditions, such as smoking and particularly obesity, have been identified as key predictors of hospitalization and critical illness, even in young adults with no underlying conditions <sup>4 5</sup>. While the pathogenesis of certain chronic diseases predisposes to severe COVID-19 outcomes, common chronic medications have been also a concern because of their potential interaction with the angiotensin-converting enzyme 2 (ACE2)<sup>6</sup>. SARS-CoV-2 binds to target cells using ACE2 in cell membranes<sup>7</sup>, an enzyme that physiologically counters the renin-angiotensin-aldosterone system (RAAS) activation, degrading angiotensin II to attenuate its subsequent physiological action. Modulation of the RAAS is a common mode of action of the widely used antihypertensive drugs ACE inhibitors (ACEI) and angiotensin II receptor blockers (ARBs) which have been reported to upregulate ACE2 expression in the heart, and mostly in animal models <sup>8-12</sup>. This potential upregulation suggested that ACE2 expression may be increased in patients treated with ACEI or ARBs, potentially worsening further the prognosis of COVID-19 infection among chronic patients, and raising early concerns during the first phases of the outbreak of the SARS-CoV-2 pandemic <sup>13 14</sup>. As patients with chronic comorbidities were also identified as more vulnerable to severe COVID-19 disease, it is necessary to understand whether part of this vulnerability could be attributed to the use of ACEI/ARBs and to evaluate the risk of discontinuing this otherwise essential, first-line therapy, for hypertensive and diabetic patients.

To date, a number of studies addressing the potential effect of ACEI and ARBs on the prognosis of COVID-19 have been reported, mostly supporting the absence of harmful effects of these drugs on COVID-19 prognosis <sup>15-32</sup>. In these studies, a wide range of statistical methods have been used to test this hypothesis, including comparison of proportions, percentage points, logistic regression, or time-to-event analysis and Cox models. The use of standard methods for these particular analyses can

1  
2  
3 100 easily lead to biased estimates, in particular when analyzing hospital data because of the presence of  
4  
5 101 competing events, such as death and recovery, and the time-dependency of these competing events  
6  
7 102 <sup>33 34</sup>. As such, the analysis of the association of ACEI/ARBs on the progression of COVID-19 or related  
8  
9  
10 103 mortality requires the assessment of competing risks/events. Analyzing time-to-event data in a multi-  
11  
12 104 state setting would better fit the true progression of COVID-19 in hospitalized patients, as shown by  
13  
14 105 two studies using a multi-state-approach in the context of COVID-19 <sup>35 36</sup>. Multi-state models allow  
15  
16 106 for studying clinically competing events (discharged alive vs deceased), as well as disease progression  
17  
18 107 (e.g. in terms of hospital stay duration, transfer to intensive care units (ICU), treatment received),  
19  
20  
21 108 simultaneously over time. This multi-state model framework ensures avoiding bias that stems from  
22  
23 109 censoring patients (informative censoring and/or selection bias) and time-dependent predictors  
24  
25 110 (time-dependent bias), as well as circumvents event-related dependency by treating disease  
26  
27 111 progression as a transient state that might influence the probability of experiencing a certain future  
28  
29 112 outcome depending on patient's risk factors. While accounting for these biases, we revisited the the  
30  
31 113 hypothesis of the potential association of ACEI/ARBs use in patient's prognosis during hospitalization  
32  
33  
34 114 using a competing risk multi-state model and nationwide hospital surveillance data on COVID-19  
35  
36 115 patients in Belgium.  
37  
38  
39 116

## 41 117 2. PATIENTS AND METHODS

### 43 118 2.1. Data sources

45 119 All methods were carried out in accordance with relevant guidelines and regulations. Nationwide  
46  
47 120 hospital surveillance data on COVID-19 patients in Belgium are routinely gathered by Sciensano, the  
48  
49 121 Belgian Institute of Public Health, which is legally entitled institution for surveillance of infectious  
50  
51 122 diseases in Belgium (Royal Decree of 21/03/2018). Retrieving informed consent was determined as a  
52  
53 123 disproportional load on the hospital resources in the crisis situation. An information letter was given  
54  
55 124 to the patients at the time of discharge which contained an explanation of their rights concerning the  
56  
57  
58 125 data that was gathered by Sciensano. The COVID-19 hospital surveillance was authorised by an  
59  
60



independent administrative authority protecting privacy and personal data and was approved by the ethical committee of Ghent University Hospital (BC-07507). Details on the Belgian COVID-19 hospital surveillance system have been previously published <sup>37</sup>. The system cover 119 hospitals in Belgium, who report standardized information on hospitalized COVID-19 patients collected through a structured questionnaire at hospital admission and discharge. An anonymized subset of data from Sciensano was shared with the Institute of Tropical Medicine through a secured data transfer platform applying data encryption. Ethical clearance was obtained from the Institutional Review Board (IRB) of the Institute of Tropical Medicine after revision of the research protocol num. 1393/20, 02/05/2020.

Variables collected at admission include the date of hospital admission, reason for hospitalization, symptoms, clinical signs, treatment with ACEI or ARBs, and demographic information such as age, gender, and the presence of chronic comorbidities. Information recorded at discharge includes laboratory values, details on COVID-19 specific treatments during hospital stay, date of discharge, health status at discharge, and measures on the severity of the disease such as the need for transfer to ICU, invasive ventilation support and/or oxygenation by extracorporeal membrane oxygenation (ECMO), and the development of a bacterial and/or fungal superinfection, pneumonia, and/or acute respiratory distress syndrome (ARDS). Dates for these severe events were only available for ICU transfer.

**2.2. Study population**

Adult COVID-19 patients with a SARS-CoV-19 infection confirmed by polymerase chain reaction (PCR), and/or suggestive imaging alterations on chest CT combined with typical clinical presentation, at admission or while hospitalized in Belgium from March 14 to June 14, 2020 were considered as COVID-19 patients (n=16,341). Of these, patients with completed questionnaires both on admission and discharge (12,109 patients, 74.1%) were selected. Information on patients admitted to hospitals before March 1 2020 (270, 1.65%) for reasons other than COVID-19, and infected while hospitalized

was also removed. Furthermore, information on patients with implausible admission dates was removed, including: Date of discharge before date of admission (42 patients, 0.25%); Date of ICU transfer preceding date of admission (31 patients, 0.19%); Date of discharge before date of ICU transfer (2 patients, 0.01%); Date of discharge preceding the date of ICU discharge if the difference was more than 1 day (47 patients, 0.28%). The final dataset for descriptive analyses included information on 11717 COVID-19 patients. For the multivariate multi-state model, patients with unknown use of ACE/ARBs (718 patients, 6.12%) were also excluded, along with those missing information on gender (118 patients, 0.72%) or unknown transfer to ICU (15, 0.09%). The final dataset for the multi-state model contained information from 103 hospitals in Belgium and 10,866 COVID-19 patients, including 539 patients (5%) that were admitted directly to ICU.

### 2.3. Study outcomes

Patients were considered to have recovered when their status at hospital discharge was recorded as “cured” or “other”. In the latter case, it was assumed they were allowed to recover at home, revalidation center or nursing home. Patients were considered as an in-hospital death when their status at discharge was recorded as “death”. Patients were considered lost to follow up when their status at discharge was recorded as “unknown” or when they were transferred to another hospital (recorded as status at discharge = “transfer to another hospital”), as no further information was available. Severe COVID-19 was captured in the database as an illness that required ECMO or artificial ventilation, or to have experienced ARDS, pneumonia, bacterial and/or fungal co-infection, or required transfer to or treatment at ICU. Among these, event date was only available for transfer to ICU, and only this variable could be therefore selected for the models as time-defined severity outcome. Incorporating the intermediate event of ICU transfer allows for a patient’s risk profile for recovery and death to be different before and after this intermediate event. In order to do so, time to severe illness was defined as the time passed from the date of hospital admission to the date of

transfer to ICU, while length of hospital stay was defined as the date from hospital admission to the date of hospital discharge (either recovery, in-hospital death, or lost to follow-up).

**2.4. Information on ACEI/ARBs and conditions related to COVID-19 prognosis**

A dedicated section in the admission dataset covered the use of ACEI, ARBs or both, without specification of the specific drug. The admission database contained information on the following factors associated with COVID-19<sup>38</sup>. Demographics (age and gender), risk factors (current smoking, high blood pressure (HBP), and obesity), prevalent comorbidities (DM, chronic renal disease, CVD, chronic lung disease, chronic liver disease, malignant solid neoplasms, hematological cancers, and immunosuppression). Smoking status was only available for 53% of the patients. Obesity also presented a large number of missing values (33.2%) because this variable was only recorded after April 3, 2020. Similarly, there were also missing values for cognitive issues (5.7%) as this variable was only recorded after March 23, 2020.

**3. STATISTICAL MODEL**

Patient’s characteristics at hospital admission, ICU stay, and at hospital discharge were visualized on histograms and summarized as means and standard deviations or counts and percentages for continuous and categorical variables, respectively. Descriptive analyses were provided for patients overall and stratified by ACEI/ARBs use, including unknown use.

To study the association of ACEI/ARBs on COVID-19 progression on a multivariate multi-state model, a first model for identifying confounders was carried out. A backwards stepwise logistic regression with ACEI/ARBs use as dependent variable, and including factors and conditions previously associated with COVID-19 outcomes and present in the database<sup>39 40</sup>, was used to inform the selection of potential confounders based on Akaike’s information criterion (AIC). The variables used in the variable selection model were: gender, age, HBP, CVD, DM, obesity, chronic renal disease, chronic liver disease, chronic lung disease, solid malignant neoplasms, hematological cancers,

immunodepression, and cognitive impairment. Two models were used depending on the availability of data, a first model (model 1) excluding variables collected at a later date (obesity, and cognitive issues) and using a full dataset (N=10,866), and a complete case analysis (model 2) excluding patients with missing data for obesity and cognitive issues (N=7,303).

We devised a multi-state model reflecting the progression of patients from hospital admission to discharge accounting for the patient's characteristics identified to be potential confounders, and introducing ACEI/ABRs use as dependent variable. The model starts with one initial state (hospitalization), a potential (transient) state defined as ICU transfer (as a proxy for severe COVID-19 disease, as only ICU transfer had an associated date in the database) and two absorbing states (in-hospital death, and recovery). The multi-state model is characterized by transition hazards between the states; defined as the instantaneous risk for moving from one health state to another. The transitions hazards are used to calculate transition probabilities, as the conditional probabilities of experiencing future outcomes, given the history and a particular set of prognostic factors (model covariates) for a given patient. The four-state model thereby comprised five possible patient's transitions; 1) hospitalization to ICU, 2) hospitalization to recovery, 3) hospitalization to in-hospital death, 4) ICU to recovery, and 5) ICU to in-hospital death, as presented in [Figure 1](#). A Cox-Markov model for the regression on the transition specific hazards was fitted using the *coxph* and *msfit* functions in R *survival* package<sup>41</sup>. This approach is equivalent of constructing five separate Cox regression models, one for each transition hazard. The cumulative baseline transition hazard (all covariate values equal to the reference value) was estimated by the Breslow estimator with the Aalen estimator of variance<sup>42</sup>. We integrated these separate Cox models in a multi-state framework studying different outcomes simultaneously and allowing the calculation of transition probabilities. The transition probabilities were then estimated by combining the baseline cumulative hazard and regression coefficients. Using R, the *mstate* package and *msfit* function was applied to obtain cumulative (baseline) transition hazards and the function *probtrans* to obtain the transition probabilities<sup>43</sup>. Estimates obtained from the Cox-Markov models are displayed in a table and

significance is established at the 5% significance level. Cumulative (baseline) transition hazard plots and transition probability plots were also generated for visual aid. In a setting with covariates, a regression model for the transition specific hazards was used, whereby the covariates may help to explain the difference in transition hazards. Model diagnostics were performed to check model assumptions of proportional hazards, linearity, and interactions. Assumptions to the Markov model were assessed by including time from hospital admission to ICU transfer in the model for transition 4 and transition 5. A relaxation of the Markov assumption was also explored in the analysis.

**4. Patient and public involvement**

As a secondary data analysis of COVID-19 surveillance data this study did not involve patients or the public in the design, conduct or dissemination plans.

**5. RESULTS**

**5.1. Descriptive analysis**

From the 11,717 patients available for this analysis, almost of all them (94.2%) presented symptoms or clinical signs compatible with COVID-19 at admission ([Supplemental Table 1](#)). Most frequent symptoms were fever (61.3%) and cough (53.2%), and most frequent signs were abnormal pulmonary imaging (63.1%) compatible with viral pneumonia, abnormal auscultation (44.8%), and dyspnea (42.4%). On admission, 15.1% of patients had a record of taking ACEI, and 8.5% ARBs, with only 0.4% taking both ACEI and ARBs. For the purpose of this analysis these patients were merged as ACEI/ARBs users. The majority of patients (69.9%) were non-users of ACEI/ARBs versus 23.9% of users, and only for a small proportion (6.1%) of patients the use of ACEI/ARBs was unknown ([Table 1](#)). No difference was seen in the frequency of signs and symptoms reported according to ACEI/ARBs use ([Supplemental Table 1](#)). Patients using ACEI/ARBs were markedly older (median [IQR] age 76 [65-84] years) than non-users (67 [53-81]) while no gender-differences were observed. As expected, ACEI/ARBs users presented more frequently (74.4%) HBP than non-users (39.2%), as well as chronic

lung disease (16.8% vs 14.4%), chronic renal disease (19.3% vs 11.1%), DM (33.3% vs 18.1%), and particularly CVD (53.1% vs 28.4%). Multiple comorbidities (HBP, DM and CVD) were more frequent among ACEI/ARBs users (15.1%) than non users (4.9%) ([Table 1](#)). During hospital stay, over 80% of COVID-19 patients experienced one severe episode of either pneumonia, superinfection, ARDS, or mechanical ventilation, and 25.9% of patients had two or more severe episodes ([Table 2](#)). The most common manifestation of COVID-19 severity was pneumonia (79.1%), followed by other infections (19.4%), ARDS (12.7%), and artificial ventilation (7.6%). Frequency of severe conditions was nearly the same for both ACEI/ARBs users and non-users (27.8% versus 25.4%). Of all admitted patients, 1,518 (13.0%) were transferred to ICU, mostly those with severe pneumonia (93.7%), or in need of artificial ventilation (58.0%), and remained at ICU for a mean duration of  $11.5 \pm 10.7$  days. Transfer to ICU was marginally more frequent among ACEI/ARBs users (15.1% versus 12.1%). Almost 78% of the patients admitted to Belgian hospitals recovered from COVID-19, either during hospitalization (51.2%) or at home or revalidation centre or nursing home (26.4%) after an average  $12.6 \pm 10.9$  days in the hospital. Only 2% of patient's information was lost to follow-up (transferred to another health care provider or unknown status at discharge).

## 5.2. Multi-state model

A multivariate state-arrival extended Cox-Markov model was used to study the potentially different progression of COVID-19 patients through health states during hospitalization according to the use of ACEI/ARBs. Possible transitions, and number of patients in each health state are represented in [Figure 1](#). The selection of variables for adjusting the models were based on backwards stepwise logistic regression of ACEI/ARBs use as a function of potential confounding factors associated with COVID-19 recorded at admission ([Table 2 Supplemental](#)). A first model (model 1) using all available patients, identified five variables associated with the use of ACEI/ARBs and COVID-19: male gender (OR 1.33, 95%CI 1.21-1.47), older age (1.01, 1.01- 1.02 per 1-year increase), and prevalent CVD (1.71, 1.55 -1.90), diabetes (1.38, 1.24-1.54), and HBP (5.65, 5.10-6.27). Additionally, a

1  
2  
3 281 second model (model 2) was used in sensitivity analysis including additional covariates (prevalent  
4  
5 282 obesity and cognitive impairment) that were only available in a subset (62%) of patients ([Table 2](#)  
6  
7 283 [Supplemental](#)). Because very few patients were asymptomatic on admission it was deemed  
8  
9  
10 284 unnecessary to adjust the regression models for severity of disease at admission. For 16 patients  
11  
12 285 (0.1%) it was unknown whether they were transferred to ICU. These patients are therefore excluded  
13  
14 286 from the multistate Cox-Markov regression analysis.

15  
16 287 Plots for the cumulative hazard and transition probability between health states considering  
17  
18 288 ACEI/ARBs use were obtained by setting all model covariates to reference values (female gender, no  
19  
20 289 CVD, no HBP, and no DM), and median (70 years) age [Figure 2](#). When looking at the cumulative  
21  
22 290 hazard for the five possible transitions ([Figure 2A](#)), the hazard for recovery was markedly greater  
23  
24 291 than that of in-hospital death. In comparison with the other cumulative hazards, the hazard for  
25  
26 292 transfer to ICU was substantially smaller, representative of most COVID-19 patients not needing  
27  
28 293 intensive care, or not meeting criteria for admission (for instance after evaluation of frailty, and  
29  
30 294 chance of survival). Transfer to ICU was associated with increased hazard for in-hospital death and  
31  
32 295 reduced hazard for recovery. The use of ACEI/ARBs was associated with a modest but significant  
33  
34 296 association with the hazard of transition 2 (more recovery) and 3 (less in-hospital death), from  
35  
36 297 admission. The use of ACEI/ARBs was not observed to be associated with transfer to ICU (transition  
37  
38 298 1), nor with recovery (transition 4) or in-hospital death (transition 5) after ICU. Overall, the  
39  
40 299 probability of being transferred to ICU was, for most patients, less than that of recovery ([Figure 2B](#)).  
41  
42 300 However, those needing ICU had a reduced probability of recovery and greater probability to  
43  
44 301 decrease in the hospital than those patients not transferred to ICU ([Figure 2C](#)).

45  
46 302 The estimates for the transition hazards for ACEI/ARBs use accounting for identified  
47  
48 303 confounding in the potential association with COVID-19 severity/fatality are presented in [Table 3](#). In  
49  
50 304 multivariate models, the use of ACE/ARBs was associated (HR 1.07, 95%CI 1.01-1.13) with more  
51  
52 305 recovery, and less death (0.83, 0.75-0.93). Even though there was a significant association between  
53  
54 306 ACEI/ARBs use the hazard of more recovery (transition 2), and less in-hospital death (transition 3)



this effect is modest, especially when reviewing state-occupation probabilities ([Figure 1 Supplemental](#)). In respect to the adjusting variables ([Table 3 Supplemental](#)), male gender and HBP were associated with transfer to ICU (severity), and older age also influenced this transition. Similarly, male gender, and older age, as well as prevalent CVD, HBP or DM were associated with less recovery. Similar to transfer to ICU, progression to death was associated with male gender and age, as well as with prevalent CVD. No other comorbidity included in this model (i.e. associated with ACEI/ARBs use) appeared to be associated with fatality. For severe patients (transferred to ICU) recovery or death depended mostly on age, albeit fatal COVID-19 was also associated with the presence of DM, and a lengthier period between admission and ICU was significantly associated with less recovery after ICU ([Table 3 Supplemental](#)). The impact of further adjustment for variables identified during confounder selection (obesity and cognitive issues) in the state transition of COVID-19 patients during hospitalization, resulted in loss of more than half of all patients due to missing values ([Table 1](#)). While estimated hazards for previous factors remained similar, the presence of cognitive issues was statistically significantly associated with transitions 1, 2, and 3 (i.e. less transfer to ICU, less recovery, and more in-hospital death), and obesity was strongly and statistically significantly associated with transition 1 only (more transfer to ICU). In this complete-case model, after additional adjustment for obesity and cognitive issues, the HR for ACEI/ARBs use for transition 2 (admission to recovery) did not remain significant probably due to a decreased statistical power, since the point estimates were similar.

## 6. DISCUSSION

In this study, a competing risk multistate model has been developed for the first time to address the magnitude and direction of the association of ACEI/ARBs use in COVID-19 prognosis. Our analyses indicate a protective association of ACEI/ARBs use, with increase recovery and survival, once important confounding factors such as age, particularly 70 and over, and male gender are accounted for. Chronic comorbidities such as CVD, HBP, and DM are also associated with less recovery in this



1  
2  
3 333 model setting. Although there is a protective association of ACEI/ARBs use on COVID-19 in-hospital  
4  
5 334 death and more recovery, this association is modest, especially when looking at the state-occupation  
6  
7 335 probabilities. In our model, once the patient progresses to a severe state, no association of  
8  
9 336 ACEI/ARBs use was observed in the transition probabilities to recovery or in-hospital death; only  
10  
11 337 older age and prevalent DM, remained significant covariates in our model, arguably because of the  
12  
13 338 smaller sample size (transfer to ICU occurred only for 13% of patients). Previous studies using the  
14  
15 339 same data source identified other comorbidities as independent risk factors for COVID-19  
16  
17 340 severity/death in ICU patients, including chronic pulmonary disease, chronic renal disease, and  
18  
19 341 immunosuppression<sup>44</sup>. Although we accounted for these factors in our model selection, they were  
20  
21 342 not selected as they are not considered to be related to the use of ACEI/ARBs but may nonetheless  
22  
23 343 constitute an independent risk factors for severity.

24  
25  
26  
27 344 Because of the important clinical relevance, there have been numerous reports on studies of  
28  
29 345 the potential association between ACEI/ARBs and (worse) prognosis of COVID-19. Early studies of  
30  
31 346 smaller sample size and mostly descriptive design pointed to either no association or moderately  
32  
33 347 lower rates of severe disease among ACEI/ARBs users<sup>15-20</sup>. Further retrospective analysis involving  
34  
35 348 larger patient samples generally reported a lack of association<sup>21</sup>. A population-based study in Italy's  
36  
37 349 Lombardy region involving 6,272 cases identified across the Regional Health Service and matched 1:5  
38  
39 350 to population-based control, found no association between the use of ACEI (adjusted OR 0.91, 95%CI  
40  
41 351 0.69-1.21) or ARBs (adjusted OR 0.83, 95%CI 0.63-1.10) and severe/fatal COVID-19<sup>22</sup>. Similarly, a  
42  
43 352 case-control study in the Spanish region of Madrid with data on 1,139 hospitalized cases matched  
44  
45 353 1:10 to population controls found no association (adjusted OR 0.94, 95%CI 0.77-1.15) of ACEI/ARBs  
46  
47 354 use and severe or fatal disease<sup>23</sup>. Analyzing data from patients in the New York University Langone  
48  
49 355 Health electronic health record who had COVID-19 test results (12,594 patients), neither an  
50  
51 356 increased likelihood of a positive test nor a severe disease status was observed for patients using  
52  
53 357 ACEI/ARBs (or any other RAAS medication) using propensity score matching<sup>24</sup>. In a nationwide study  
54  
55 358 in Korea using insurance claims of 66,793 individuals tested for COVID-19, the use of ACEI/ARBs was  
56  
57  
58  
59  
60

not associated with a higher risk of mortality (adjusted OR 0.88, 95%CI 0.53-1.44)<sup>25</sup>. Similarly, a large retrospective analysis of an Italian registry cohort including 43,000 patients concluded that neither ACEI (adjusted HR 0.97, 95%CI 0.89-1.06) or ARB (adjusted HR 0.98, 95% CI 0.89-1.06) use was associated with either an increased or decreased risk of all-cause mortality<sup>26</sup>. A multicenter study with 1,128 hypertensive patients, and using mixed-effect Cox models (site as a random effect, and model adjusted for age, gender, comorbidities, and in-hospital medications) reported a lower risk for all-cause mortality in the ACEI/ARB patients versus the non-ACEI/ARB group (adjusted HR 0.42, 95%CI 0.19-0.92), and further compared with the use of other antihypertensive drugs, (adjusted HR 0.30, 95%CI, 0.12-0.70)<sup>27</sup>. Previous studies using Cox models reported also a reduced mortality risk for patients using ACEI/ARBs<sup>27 28</sup>. In others, albeit not statistically significant, estimates were very similar to the ones reported in our study for mortality (adjusted HR 0.83, 95%CI 0.67-1.03) and for severe disease (adjusted HR 1.15, 95%CI 0.95-1.41)<sup>29</sup>. Similarly, but outside of the hospital setting, studies with data from general practitioners in England, found a strong association of ACEI/ARBs use and a reduced risk of COVID-19 disease (HR 0.63, 95%CI 0.59-0.67) albeit not severity (HR 1.02, 95%CI 0.83-1.25), and marked interactions with ethnicity with higher risks observed for Black Africans compared to Whites<sup>30</sup>. Variations observed between different ethnicities raise the possibility of specific associations of ACEI/ARBs on COVID-19 disease susceptibility and severity which deserves further study. Furthermore, three review papers on the topic have concluded there is either no difference or a reduced risk when looking at mortality and/or severe disease<sup>21 31 32</sup>, and no evidence to support discontinuing the treatment with ACEI/ARBs<sup>45</sup>. This substantial body of evidence seems aligned with recent findings from clinical studies that do not support the hypothesis of an increased expression of ACE2 in chronic patients treated with ACEI or ARBs as a driver of severe COVID-19<sup>10-12</sup>.

Whereas the results on an increased risk of severe/fatal COVID-19 in association with the outpatient use of ACEIs/ARBs appear to point in the same direction, studies on the potential role of in-hospital use of ACEI/ARBs have described a protective association of continuing the treatment throughout hospitalization<sup>27 46 47</sup>. In a multi-center study including 1,128 adult patients with HTN and

1  
2  
3 385 diagnosed with COVID-19, including 188 taking ACEI/ARB and 940 without using ACEI/ARB during  
4  
5 386 hospitalization, the risk for all-cause mortality was lower in the ACEI/ARB group versus the non-  
6  
7 387 ACEI/ARB group (HR 0.42, 95% CI, 0.19–0.92) <sup>27</sup>. Similarly, a study conducted in 347 patients  
8  
9 388 hospitalized for COVID-19 in Paris (France) analysing the association between in-hospital exposure of  
10  
11 389 ACEI/ARBs and mortality within 30 days of hospital admission using logistic regression analysis, no  
12  
13 390 association (OR 0.62, 95%CI: 0.25-1.48) based on chronic exposure but a protective association (OR  
14  
15 391 0.25, 95%CI: 0.09-0.65) based on in-hospital exposure was observed.<sup>46</sup> Among 397 patients with  
16  
17 392 COVID-19 addmitted to hospitals in Rozzano-Milan (Italy) the risk of mortality was significantly  
18  
19 393 reduced in patients who continued ACEI/ARBs as compared with those who discontinued and those  
20  
21 394 not taking ACEI/ARB therapy (OR 0.14, [95%CI: 0.03-0.66]) <sup>47</sup>. Using data from 7 hospitals in Madrid  
22  
23 395 (Spain), no difference in mortality rates was observed among patients that discontinued (340  
24  
25 396 patients) ACEI/ARBs treatment (HR 1.01, [95%CI 0.70–1.46]) versus those that continued (280  
26  
27 397 patients) <sup>48</sup>. Furthermore, recent data from two randomized trials could not confirm any impact on  
28  
29 398 clinical outcomes in hospitalized COVID-19 patients discontinuing treatment of ACEI or ARBs as  
30  
31 399 compared with those continuing their treatment <sup>49 50</sup>, hereby further supporting a safe ACEI/ARBS  
32  
33 400 treatment continuation if there is an indication for treatment.  
34  
35  
36  
37  
38  
39  
40

41 402 Our study builds on these previous reports where standard statistical models were used for  
42  
43 403 analysis, introducing a model framework overcoming the risk of biases <sup>33</sup>. Logistic-based regression  
44  
45 404 models might introduce selection bias by excluding patients who are still hospitalized at the last  
46  
47 405 follow-up, hence the need for time-to-event models that allow patient censoring <sup>33</sup>. These time-to-  
48  
49 406 event models, such as Cox regression models, should preferably account for the presence of  
50  
51 407 competing risks to avoid informative censoring bias, and for time-dependent predictors to  
52  
53 408 appropriately handle index time or follow-up time of covariates<sup>33 49</sup>. Integrating standard Cox models  
54  
55 409 into a multi-state framework allows the study of separate outcomes simultaneously and allows the  
56  
57 410 calculation of the transition probabilities, adding a layer of interpretation. In this way, by  
58  
59  
60

incorporating event-related dependency, i.e., transitions to intermediate events that likely influence disease progression, the multi-state model more accurately describes the evolution of COVID-19 in hospitalized patients<sup>51</sup>. In the present study, we used a time-to-event analysis in a multi-state framework considering competing risks to account appropriately for censoring<sup>52</sup>, thereby robustly showing a modest, yet significant, positive association of ACEI/ARBs use in recovery and survival of hospitalized COVID-19 patients accounting for confounding factors.

Our study uses nationwide hospital surveillance data, with mandatory participation, and includes all general hospitals (including university hospitals) in Belgium, both those managed by a public authority and privately managed are represented. The surveillance does not cover psychiatric hospitals or specialist hospitals<sup>37</sup>. The use of comprehensive datasets, but more so the utilization of models adequately fitting to time-to-event hospital data with mutually exclusive health states results in less probability of introducing biases and are crucial for correct evidence-based information for decision making. Our study makes some assumptions, and unknowns such as the lack of information on ACEI/ARBs exact indication and whether their use was continued after admission. This lack of accounting for time-varying exposure introduces a risk of immeasurable time bias<sup>34</sup>, as seen in many reports, though assumed to be minimal because no difference in COVID-19 severity outcomes between treatment discontinuation vs continuation<sup>49 50</sup>. Our models are not adjusted for severity at baseline since we reasoned that hospital admission was already an indicator of severe disease and 94% of patients had symptoms compatible with COVID-19 diagnosis at baseline. Further, even though other events potentially indicating severity (ECMO, ARDS, pneumonia, bacterial and/or fungal co-infection) were available in the database, only transfer to ICU was linked to a calendar date and was therefore the only event which could be used as a proxy for severe health state in our time-dependent model, indicating that our estimates might represent more a critical state of the patient. In addition, admission to intensive care is not solely based on the clinical status of the patient, but also on other criteria such as frailty. Also, ICU admission criteria might have been more restrictive in

1  
2  
3 437 the peak period of the epidemic whilst certain ICU were overloaded. Because the surveillance data is  
4  
5 438 limited to the most important variables, we cannot discard the possibility of some degree of residual  
6  
7 439 confounding in our results. An important limitation of our main analysis is the impossibility of  
8  
9 440 adjusting our models for smoking status, obesity and cognitive issues at baseline. Using available  
10  
11 441 smoking information was not deemed appropriate due to the excessive number of missing values,  
12  
13 442 and the lack of information of the reason for the incomplete data. We used, however, data on  
14  
15 443 obesity and cognitive issues, which collection was introduced later, in a complete case analysis to  
16  
17 444 confirm the results obtained in the main model. Nevertheless, these analyses on a reduced sample of  
18  
19 445 patients should be interpreted with caution as a time effect is likely present because of the late data  
20  
21 446 collection. Finally, our analyses are based on patient's medical files and rely on how clinicians  
22  
23 447 reported clinical observations and anamnesis which might vary across hospitals, and are  
24  
25 448 representative of the first so called wave of the epidemic in Belgium, and associations might differ in  
26  
27 449 subsequent studies and in other settings.  
28  
29  
30  
31  
32  
33

34 451 **7. CONCLUSIONS**

35  
36 452  
37  
38 453 After adjustment for important confounders there is modest, yet significant, positive association  
39  
40 454 of ACEI/ARBs use on recovery and survival of hospitalized COVID-19 patients, without affecting  
41  
42 455 admission to intensive care. This supports the use of ACEI/ARB in those patients who need them, also  
43  
44 456 when needing hospitalization from COVID-19. These findings are based on an analytical model that  
45  
46 457 adequately fits hospital data, where patients progress across different, competing, health states  
47  
48 458 providing a more complete and accurate view of the research question within a reduced risk of bias  
49  
50 459 framework. Integrating standard cox models into a multi-state framework allows the study of  
51  
52 460 separate outcomes simultaneously and allows the calculation of the transition probabilities, adding a  
53  
54 461 layer of interpretation. Multi-state models should be favoured over separate survival analysis when  
55  
56  
57  
58  
59  
60

competing risks are present, and traditional methods such as logit functions should be discouraged when time-to-event is available.

#### **Contributorship statement**

JLP, MvdS, MAW, DV conceptualized the study. DV performed data curation and provided data. JLP, EG, EM, DS developed methodology, and performed analysis. JLP supervised the study, and drafted the manuscript. All authors have critically reviewed, commented, and approved the manuscript before submission.

#### **Competing interest**

The authors declare that they have no competing interests.

#### **Funding statement**

This research received no specific grant from any funding agency in the public, commercial or not-for-profit sectors.

#### **Data sharing statement**

The data that support the findings of this study are available from Sciensano but restrictions apply to the availability of these data, which were used under license for the current study, and so are not publicly available. Data are however available from the authors upon reasonable request and with permission of Sciensano.

#### **Acknowledgments**

The authors thank the Members of the Belgian Collaborative Group on COVID-19 Hospital Surveillance: Samy Amir Aouachria, Centre Hospitalier Chrétien, Liège, Belgium; Kristof Bafort, Mariaziekenhuis, Pelt, Belgium; Leïla Belkhir, Cliniques Universitaires Saint-Luc, Brussels, Belgium;

1  
2  
3 488 Koen Blot, Sciensano, Belgium; Nathalie Bossuyt, Sciensano, Belgium; Vincent Colombie, Centre  
4  
5 489 Hospitalier Epicura, Baudour, Belgium; Nicolas Dauby, Centre Hospitalier Universitaire Saint-Pierre,  
6  
7 490 Brussels, Belgium; Paul De Munter , Universitair Ziekenhuis Leuven, Leuven, Belgium; Robby De  
8  
9 491 Pauw, Sciensano, Brussels, Belgium; Didier Delmarcelle, Clinique St. Jean, Brussels, Belgium; Mélanie  
10  
11 492 Delvallee, Centre Hospitalier de Wallonie Picarde, Tournai, Belgium; Rémy Demeester, Centre  
12  
13 493 Hospitalier Universitaire de Charleroi, Charleroi, Belgium; Thierry Dugernier, Clinique Saint-Pierre,  
14  
15 494 Ottignies, Belgium; Xavier Holemans, Grand Hôpital de Charleroi, Charleroi, Belgium; Benjamin  
16  
17 495 Kerzmann, Clinique Notre Dame de Grâce, Gosselies, Belgium; Pierre Yves Machurot, Centre  
18  
19 496 Hospitalier de l'Ardenne, Belgium; Philippe Minette, Centres Hospitaliers Jolimont, Belgium; Jean-  
20  
21 497 Marc Minon, Centre Hospitalier Régional de la Citadelle, Liège, Belgium; Saphia Mokrane, Hôpitaux  
22  
23 498 Iris Sud, Brussels, Belgium; Catherine Nachtergal, Cliniques de l'Europe, Brussels, Belgium; Séverine  
24  
25 499 Noirhomme, Centre Hospitalier Régional de Namur, Belgium; Denis Piérard, Universitair Ziekenhuis  
26  
27 500 Brussel, Brussels; Camelia Rossi, Centre Hospitalier Universitaire Ambroise Paré, Mons, Belgium;  
28  
29 501 Carole Schirvel, CHIREC, Brussels, Belgium; Erica Sermijn, A.S.Z. Ziekenhuis, Aalst, Belgium; Ben  
30  
31 502 Serrien, Sciensano, Belgium; Fabio Taccone, Hôpital Erasme, Brussels, Belgium; Frank Staelens, OLV  
32  
33 503 Ziekenhuis, Aalst, Belgium; Filip Triest, Algemeen Ziekenhuis Sint Lucas, Gent, Belgium; Nina Van  
34  
35 504 Goethem, Sciensano, Brussels, Belgium; Jens Van Praet, Algemeen Ziekenhuis Sint Jan, Brugge-  
36  
37 505 Oostende, Belgium; Anke Vanhoenacker, Ziekenhuisnetwerk Antwerpen, Belgium; Roeland  
38  
39 506 Verstraete, Algemeen Ziekenhuis, Monica, Belgium; Elise Willems, Algemeen Ziekenhuis Nikolaas,  
40  
41 507 Sint-Niklaas, Belgium; Chloé Wyndham-Thomas, Sciensano, Belgium.  
42  
43 508  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

509

For peer review only



1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

510   **REFERENCES**

512   1. Yang X, Yu Y, Xu J, et al. Clinical course and outcomes of critically ill patients with SARS-CoV-2  
513       pneumonia in Wuhan, China: a single-centered, retrospective, observational study. *Lancet*  
514       *Respir Med* 2020;8(5):P485-P81. doi: 10.1016/S2213-2600(20)30079-5 [published Online  
515       First: 2020/02/28]

516   2. Guan WJ, Ni ZY, Hu Y, et al. Clinical Characteristics of Coronavirus Disease 2019 in China. *N Engl J*  
517       *Med* 2020;382(18):1708-30. doi: 10.1056/NEJMoa2002032 [published Online First:  
518       2020/02/29]

519   3. Zhang JJ, Dong X, Cao YY, et al. Clinical characteristics of 140 patients infected with SARS-CoV-2 in  
520       Wuhan, China. *Allergy* 2020;75(7):1730-41. doi: 10.1111/all.14238 [published Online First:  
521       2020/02/23]

522   4. Kassir R. Risk of COVID-19 for patients with obesity. *Obesity Reviews*;21(6):e13034. doi:  
523       10.1111/obr.13034

524   5. Petrilli CM, Jones SA, Yang J, et al. Factors associated with hospital admission and critical illness  
525       among 5279 people with coronavirus disease 2019 in New York City: prospective cohort  
526       study. *BMJ* 2020;360:m1966. doi: 10.1101/2020.04.08.20057794

527   6. Wan Y, Shang J, Graham R, et al. Receptor Recognition by the Novel Coronavirus from Wuhan: an  
528       Analysis Based on Decade-Long Structural Studies of SARS Coronavirus. *Journal of Virology*  
529       2020;94(7):e00127-20. doi: 10.1128/jvi.00127-20

530   7. Rossi GP, Sanga V, Barton M. Potential harmful effects of discontinuing ACE-inhibitors and ARBs in  
531       COVID-19 patients. *Elife* 2020;9:e57278. doi: 10.7554/eLife.57278

532   8. Ishiyama Y, Gallagher PE, Averill DB, et al. Upregulation of angiotensin-converting enzyme 2 after  
533       myocardial infarction by blockade of angiotensin II receptors. *Hypertension* 2004;43(5):970-  
534       6. doi: 10.1161/01.HYP.0000124667.34652.1a [published Online First: 2004/03/10]

9. Patel AB, Verma A. COVID-19 and Angiotensin-Converting Enzyme Inhibitors and Angiotensin Receptor Blockers: What Is the Evidence? *JAMA* 2020;323(18):1769-70. doi: 10.1001/jama.2020.4812
10. Emilsson V, Gudmundsson EF, Aspelund T, et al. Antihypertensive medication uses and serum ACE2 levels. *medRxiv* 2020:2020.05.21.20108738. doi: 10.1101/2020.05.21.20108738
11. Jiang X, Eales JM, Scannali D, et al. Hypertension and renin-angiotensin system blockers are not associated with expression of angiotensin-converting enzyme 2 (ACE2) in the kidney. *European Heart Journal* 2020;41(48):4580-88. doi: 10.1093/eurheartj/ehaa794
12. Sama IE, Ravera A, Santema BT, et al. Circulating plasma concentrations of angiotensin-converting enzyme 2 in men and women with heart failure and effects of renin-angiotensin-aldosterone inhibitors. *European Heart Journal* 2020;41(19):1810-17. doi: 10.1093/eurheartj/ehaa373
13. Zheng Y-Y, Ma Y-T, Zhang J-Y, et al. COVID-19 and the cardiovascular system. *Nature Reviews Cardiology* 2020 doi: 10.1038/s41569-020-0360-5
14. Fang L, Karakiulakis G, Roth M. Are patients with hypertension and diabetes mellitus at increased risk for COVID-19 infection? *Lancet Respir Med* 2020;8(4):e21. doi: 10.1016/S2213-2600(20)30116-8 [published Online First: 2020/03/15]
15. Peng YD, Meng K, Guan HQ, et al. [Clinical characteristics and outcomes of 112 cardiovascular disease patients infected by 2019-nCoV]. *Zhonghua Xin Xue Guan Bing Za Zhi* 2020;48(6):450-55. doi: 10.3760/cma.j.cn112148-20200220-00105 [published Online First: 2020/03/04]
16. Huang Z, Cao J, Yao Y, et al. The effect of RAS blockers on the clinical characteristics of COVID-19 patients with hypertension. *Ann Transl Med* 2020;8(7):430. doi: 10.21037/atm.2020.03.229 [published Online First: 2020/05/13]
17. Li J, Wang X, Chen J, et al. Association of Renin-Angiotensin System Inhibitors With Severity or Risk of Death in Patients With Hypertension Hospitalized for Coronavirus Disease 2019

1  
2  
3 560 (COVID-19) Infection in Wuhan, China. *JAMA Cardiol* 2020;5(7):825-30. doi:  
4  
5 561 10.1001/jamacardio.2020.1624 [published Online First: 2020/04/24]  
6  
7 562 18. Meng X, Liu Y, Wei C, et al. Angiotensin converting enzyme inhibitors and angiotensin receptor  
8  
9 563 blockers improved the outcome of patients with severe COVID-19 and hypertension. *Sci*  
10  
11 564 *China Life Sci* 2020;1-4 doi: 10.1007/s11427-020-1813-0 [published Online First: 2020/09/26]  
12  
13  
14 565 19. Feng Y, Ling Y, Bai T, et al. COVID-19 with Different Severities: A Multicenter Study of Clinical  
15  
16 566 Features. *Am J Respir Crit Care Med* 2020;201(11):1380-88. doi: 10.1164/rccm.202002-  
17  
18 567 0445OC [published Online First: 2020/04/11]  
19  
20  
21 568 20. Yang G, Tan Z, Zhou L, et al. Effects of Angiotensin II Receptor Blockers and ACE (Angiotensin-  
22  
23 569 Converting Enzyme) Inhibitors on Virus Infection, Inflammatory Status, and Clinical Outcomes  
24  
25 570 in Patients With COVID-19 and Hypertension: A Single-Center Retrospective Study.  
26  
27 571 *Hypertension* 2020;76(1):51-58. doi: 10.1161/HYPERTENSIONAHA.120.15143 [published  
28  
29 572 Online First: 2020/04/30]  
30  
31  
32 573 21. Zhang J, Wang M, Ding W, et al. The interaction of RAAS inhibitors with COVID-19: Current  
33  
34 574 progress, perspective and future. *Life Sci* 2020;257:118142. doi: 10.1016/j.lfs.2020.118142  
35  
36 575 [published Online First: 2020/07/28]  
37  
38  
39 576 22. Mancia G, Rea F, Ludergnani M, et al. Renin-Angiotensin-Aldosterone System Blockers and the  
40  
41 577 Risk of Covid-19. *N Engl J Med* 2020;382(25):2431-40. doi: 10.1056/NEJMoa2006923  
42  
43 578 [published Online First: 2020/05/02]  
44  
45  
46 579 23. de Abajo FJ, Rodriguez-Martin S, Lerma V, et al. Use of renin-angiotensin-aldosterone system  
47  
48 580 inhibitors and risk of COVID-19 requiring admission to hospital: a case-population study.  
49  
50 581 *Lancet* 2020;395(10238):1705-14. doi: 10.1016/S0140-6736(20)31030-8 [published Online  
51  
52 582 First: 2020/05/18]  
53  
54  
55 583 24. Reynolds HR, Adhikari S, Pulgarin C, et al. Renin-Angiotensin-Aldosterone System Inhibitors and  
56  
57 584 Risk of Covid-19. *N Engl J Med* 2020;382(25):2441-48. doi: 10.1056/NEJMoa2008975  
58  
59 585 [published Online First: 2020/05/02]  
60

25. Jung SY, Choi JC, You SH, et al. Association of renin-angiotensin-aldosterone system inhibitors with COVID-19-related outcomes in Korea: a nationwide population-based cohort study. *Clin Infect Dis* 2020;71(16):2121–28. doi: 10.1093/cid/ciaa624 [published Online First: 2020/05/23]
26. Trifiro G, Massari M, Da Cas R, et al. Renin-Angiotensin-Aldosterone System Inhibitors and Risk of Death in Patients Hospitalised with COVID-19: A Retrospective Italian Cohort Study of 43,000 Patients. *Drug Saf* 2020;1-12 doi: 10.1007/s40264-020-00994-5 [published Online First: 2020/08/28]
27. Zhang P, Zhu L, Cai J, et al. Association of Inpatient Use of Angiotensin-Converting Enzyme Inhibitors and Angiotensin II Receptor Blockers With Mortality Among Patients With Hypertension Hospitalized With COVID-19. *Circulation Research* 2020;126(12):1671-81. doi: 10.1161/CIRCRESAHA.120.317134 [published Online First: 2020/04/18]
28. Gao C, Cai Y, Zhang K, et al. Association of hypertension and antihypertensive treatment with COVID-19 mortality: a retrospective observational study. *Eur Heart J* 2020;41(22):2058-66. doi: 10.1093/eurheartj/ehaa433 [published Online First: 2020/06/05]
29. Fosbol EL, Butt JH, Ostergaard L, et al. Association of Angiotensin-Converting Enzyme Inhibitor or Angiotensin Receptor Blocker Use With COVID-19 Diagnosis and Mortality. *JAMA* 2020;324(2):168-77. doi: 10.1001/jama.2020.11301 [published Online First: 2020/06/20]
30. Hippisley-Cox J, Young D, Coupland C, et al. Risk of severe COVID-19 disease with ACE inhibitors and angiotensin receptor blockers: cohort study including 8.3 million people. *Heart* 2020;106(19):1503-11. doi: 10.1136/heartjnl-2020-317393 [published Online First: 2020/08/02]
31. Fernandez-Ruiz I. RAAS inhibitors do not increase the risk of COVID-19. *Nat Rev Cardiol* 2020;17(7):383. doi: 10.1038/s41569-020-0401-0 [published Online First: 2020/05/24]

32. Jarcho JA, Ingelfinger JR, Hamel MB, et al. Inhibitors of the Renin-Angiotensin-Aldosterone System and Covid-19. *N Engl J Med* 2020;382(25):2462-64. doi: 10.1056/NEJMe2012924 [published Online First: 2020/05/02]
33. Wolkewitz M, Lambert J, von Cube M, et al. Statistical Analysis of Clinical COVID-19 Data: A Concise Overview of Lessons Learned, Common Errors and How to Avoid Them. *Clin Epidemiol* 2020;12:925-28. doi: 10.2147/CLEP.S256735 [published Online First: 2020/09/19]
34. Cohen JB, D'Agostino McGowan L, Jensen ET, et al. Evaluating sources of bias in observational studies of angiotensin-converting enzyme inhibitor/angiotensin II receptor blocker use during COVID-19: beyond confounding. *Journal of hypertension* 2021;39(4):795-805. doi: 10.1097/HJH.0000000000002706
35. Hazard D, Kaier K, von Cube M, et al. Joint analysis of duration of ventilation, length of intensive care, and mortality of COVID-19 patients: a multistate approach. *BMC Medical Research Methodology* 2020;20(1):206. doi: 10.1186/s12874-020-01082-z
36. von Cube M, Grodd M, Wolkewitz M, et al. Harmonizing Heterogeneous Endpoints in Coronavirus Disease 2019 Trials Without Loss of Information. *Critical Care Medicine* 2020;49(1):e11-e19. doi: 10.1101/2020.03.31.20049007
37. Van Goethem N, Vilain A, Wyndham-Thomas C, et al. Rapid establishment of a national surveillance of COVID-19 hospitalizations in Belgium. *Arch Public Health* 2020;78(1):121. doi: 10.1186/s13690-020-00505-z [published Online First: 2020/12/10]
38. WHO. COVID-19 and NCDs 2020 [Available from: <https://www.who.int/internal-publications-detail/covid-19-and-ncds> accessed March 27 2020.
39. de Azambuja E, Brandão M, Wildiers H, et al. Impact of solid cancer on in-hospital mortality overall and among different subgroups of patients with COVID-19: a nationwide, population-based analysis. *ESMO Open* 2020;5(5):e000947. doi: 10.1136/esmoopen-2020-000947
40. Catteau L, Dauby N, Montourcy M, et al. Low-dose hydroxychloroquine therapy and mortality in hospitalised patients with COVID-19: a nationwide observational study of 8075 participants.

- 636 *International Journal of Antimicrobial Agents* 2020;56(4):106144. doi:
- 637 <https://doi.org/10.1016/j.ijantimicag.2020.106144>
- 638 41. Survival R package. [Available from: [https://cran.r-](https://cran.r-project.org/web/packages/survival/survival.pdf)
- 639 [project.org/web/packages/survival/survival.pdf](https://cran.r-project.org/web/packages/survival/survival.pdf) accessed 1 November 2020.
- 640 42. de Wreede LC, Fiocco M, Putter H. The mstate package for estimation and prediction in non- and
- 641 semi-parametric multi-state and competing risks models. *Comput Methods Programs Biomed*
- 642 2010;99(3):261-74. doi: 10.1016/j.cmpb.2010.01.001 [published Online First: 2010/03/17]
- 643 43. Mstate R package. [Available from: <https://cran.r-project.org/web/packages/mstate/mstate.pdf>
- 644 accessed 1 November 2020.
- 645 44. Taccone FS, Vangoethem N, Depauw R, et al. The role of organizational characteristics on the
- 646 outcome of COVID-19 patients admitted to the ICU in Belgium. *The Lancet Regional Health -*
- 647 *Europe* 2020;0(0):100019. doi: <https://doi.org/10.1016/j.lanepe.2020.100019>
- 648 45. Danser AHJ, Epstein M, Batlle D. Renin-Angiotensin System Blockers and the COVID-19 Pandemic:
- 649 At Present There Is No Evidence to Abandon Renin-Angiotensin System Blockers.
- 650 *Hypertension* 2020;75(6):1382-85. doi: 10.1161/hypertensionaha.120.15082 [published
- 651 Online First: 2020/03/27]
- 652 46. Lahens A, Mullaert J, Gressens S, et al. Association between renin-angiotensin-aldosterone
- 653 system blockers and outcome in coronavirus disease 2019: analysing in-hospital exposure
- 654 generates a biased seemingly protective effect of treatment. *J Hypertens* 2021;39(2):367-75.
- 655 doi: 10.1097/hjh.0000000000002658 [published Online First: 2020/10/07]
- 656 47. Cannata F, Chiarito M, Reimers B, et al. Continuation versus discontinuation of ACE inhibitors or
- 657 angiotensin II receptor blockers in COVID-19: effects on blood pressure control and mortality.
- 658 *European Heart Journal - Cardiovascular Pharmacotherapy* 2020;6(6):412-14. doi:
- 659 10.1093/ehjcvp/pvaa056
- 660 48. de Abajo FJ, Rodríguez-Miguel A, Rodríguez-Martín S, et al. Impact of in-hospital discontinuation
- 661 with angiotensin receptor blockers or converting enzyme inhibitors on mortality of COVID-19

1  
2  
3 662 patients: a retrospective cohort study. *BMC Medicine* 2021;19(1):118. doi: 10.1186/s12916-  
4  
5 663 021-01992-9  
6  
7 664 49. Cohen JB, Hanff TC, William P, et al. Continuation versus discontinuation of renin–angiotensin  
8  
9  
10 665 system inhibitors in patients admitted to hospital with COVID-19: a prospective, randomised,  
11  
12 666 open-label trial. *The Lancet Respiratory Medicine* 2021;9(3):275-84. doi:  
13  
14 667 [https://doi.org/10.1016/S2213-2600\(20\)30558-0](https://doi.org/10.1016/S2213-2600(20)30558-0)  
15  
16 668 50. Lopes RD, Macedo AVS, de Barros ESPGM, et al. Effect of Discontinuing vs Continuing  
17  
18 669 Angiotensin-Converting Enzyme Inhibitors and Angiotensin II Receptor Blockers on Days Alive  
19  
20 670 and Out of the Hospital in Patients Admitted With COVID-19: A Randomized Clinical Trial.  
21  
22  
23 671 *Jama* 2021;325(3):254-64. doi: 10.1001/jama.2020.25864 [published Online First:  
24  
25 672 2021/01/20]  
26  
27 673 51. Meira-Machado L, de Uña-Alvarez J, Cadarso-Suárez C, et al. Multi-state models for the analysis  
28  
29 674 of time-to-event data. *Stat Methods Med Res* 2009;18(2):195-222. doi:  
30  
31 675 10.1177/0962280208092301 [published Online First: 2008/06/18]  
32  
33 676 52. Putter H, Fiocco M, Geskus RB. Tutorial in biostatistics: competing risks and multi-state models.  
34  
35 677 *Stat Med* 2007;26(11):2389-430. doi: 10.1002/sim.2712 [published Online First: 2006/10/13]  
36  
37 678  
38  
39 679  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

**FIGURE TITLES**

**Figure 1.** Schematic representation of the competing risk multi-state model and transition event matrix (number (%) patients in each transition).

*Figure footnote: Numbers in superscript represent transitions depicted in the figure. \*539 patients were directly admitted to ICU*

**Figure 2.** Plots for cumulative transition hazards (A), and state transition probabilities (B), and transition probabilities after transfer to intensive care (C) in a multi-state competing risk model considering ACEI/ARBs use (dashed line) versus no use (solid line).

**Figure 1 Supplemental.** Staked probability plot of the state-occupation probabilities in a multi-state competing risk model considering ACEI/ARBs use (A) versus no use (B).



**Table 1.** COVID-19 patient’s characteristics at hospital admission according to ACEI/ARBs use

	Total	ACEI/ARBs		
	(n = 11717)	No use (n = 8189, 69.9%)	Use (n = 2810, 23.9%)	Unknown use (n = 718, 6.1%)
<b>Demographics</b>				
Age (years) (mean (SD))	67.82 (17.17)	65.70 (17.90)	74.08 (12.85)	67.47 (17.50)
Equal or more than 70 years old (n, %)	6044 (51.6)	3791 (46.3)	1886 (67.1)	367 (51.1)
Sex (n, % males)	6154 (52.5)	4227 (51.6)	1562 (55.6)	365 (50.8)
Missing (n, %)	129 (1.1)	93 (1.1)	25 (0.9)	11 (1.5)
<b>Risk factors</b>				
Smokers (n, %)	606 (5.2)	440 (5.4)	142 (5.1)	24 (3.3)
Missing (n, %)	5413 (46.2)	3667 (44.8))	1160 (41.3)	586 (81.6)
Flu vaccination (n, %)	841 (7.2)	572 (7.0)	250 (8.9)	19 (2.6)
Missing (n, %)	10076 (86.0)	7018 (85.7)	2374 (84.5)	684 (95.3)
Obesity (n, %)*	782 (6.7)	478 (5.8)	271 (9.6)	33 (4.6)
Missing (n, %)	3887 (33.2)	2735 (33.4)	870 (31.0)	282 (39.3)
<b>Chronic comorbidities</b>				
HBP (n, %)	4593 (39.2)	2343 (28.6)	2090 (74.4)	160 (22.3)
DM (n, %)	2522 (21.5)	1486 (18.1)	936 (33.3)	100 (13.9)
Chronic renal disease (n, %)	1513 (12.9)	911 (11.1)	541 (19.3)	61 (8.5)
CVD (n, %)	3984 (34.0)	2326 (28.4)	1493 (53.1)	165 (23.0)
Chronic lung disease (n, %)	1731 (14.8)	1180 (14.4)	473 (16.8)	78 (10.9)
Cognitive impairment (n, %)**	1320 (11.3)	922 (11.3)	331 (11.8)	67 (9.3)
Missing (n, %)	668 (5.7)	461 (5.6)	173 (6.2)	34 (4.7)
Chronic neuro-muscular disease (n, %)	993 (8.5)	704 (8.6)	241 (8.6)	48 (6.7)
Solid malignant neoplasms (n, %)	990 (8.4)	697 (8.5)	261 (9.3)	32 (4.5)
Chronic liver disease (n, %)	301 (2.6)	210 (2.6)	79 (2.8)	12 (1.7)
Immunodepression (n, %)	297 (2.5)	224 (2.7)	64 (2.3)	9 (1.3)
Hematological cancers (n, %)	216 (1.8)	154 (1.9)	56 (2.0)	6 (0.8)
<b>Combination of comorbidities</b>				
None (n, %)	4760 (40.6)	4145 (50.6)	192 (6.8)	423 (58.9)
CVD & HBP (n, %)	1386 (11.8)	713 (8.7)	633 (22.5)	41 (5.7)
CVD & DM (n, %)	385 (3.3)	248 (3.0)	113 (4.0)	24 (3.3)
HBP & DM (n, %)	682 (5.8)	348 (4.2)	309 (11.0)	25 (3.5)
CVD & HBP & DM (n, %)	401 (50.6)	401 (4.9)	423 (15.1)	20 (2.8)

ACEI/ARBs, Angiotensin converting enzyme inhibitors / Angiotensin receptor blockers; CVD, cardiovascular disease; DM, diabetes mellitus; HBP, high blood pressure. \*Values only collected after April 3, 2020. \*\*Values only reported after March 23, 2020.

**Table 2.** Frequency of severity events among COVID-19 patients during hospital stay and recorded outcomes at discharge according to ACEI/ARBs use at admission

		ACEI/ARBs			
		Total (n = 11717)	No use (n = 8189, 69.9%)	Use (n = 2810, 23.9%)	Unknown use (n = 718, 6.1%)
Severe conditions					
Pneumonia (n, %)		9265 (79.1)	6501 (79.4)	2260 (80.4)	504 (70.2)
Missing (n, %)		532 (4.5)	372 (4.5)	73 (73)	87 (87)
Superinfection (n, %)		2268 (19.4)	1548 (18.9)	589 (21.0)	131 (18.2)
Missing (n, %)		1277 (10.9)	820 (10.0)	320 (11.4)	137 (19.1)
ARDS (n, %)		1492 (12.7)	996 (12.2)	389 (13.8)	107 (14.9)
Missing (n, %)		1047 (8.9)	671 (8.2)	263 (9.4)	113 (15.7)
Mechanical ventilation (n, %)		893 (7.6)	571 (7.0)	249 (8.9)	73 (10.2)
Missing (n, %)		627 (5.4)	383 (4.7)	170 (6.0)	74 (10.3)
Number of severe conditions (n, %)					
	None	2143 (18.3)	1492 (18.2)	465 (16.5)	186 (25.9)
	One	6537 (55.8)	4620 (56.4)	1565 (55.7)	352 (49.0)
	Two or more	3037 (25.9)	2077 (25.4)	780 (27.8)	180 (25.1)
Intensive care					
Transfer to ICU (n, %)		1518 (13.0)	990 (12.1)	425 (15.1)	103 (14.3)
Missing (n, %)		16 (0.1)	16 (0.2)	0 (0)	0 (0)
Transfer to ICU + pneumonia (n, %)		1423 (93.7)	932 (94.1)	395 (92.9)	96 (93.2)
Transfer to ICU + superinfection (n, %)		653 (43.0)	423 (42.7)	181 (42.6)	49 (47.6)
Transfer to ICU + ARDs (n, %)		831 (54.7)	547 (55.3)	224 (52.7)	60 (58.3)
Transfer to ICU + mechanical ventilation (n, %)		880 (58.0)	561 (56.7)	246 (57.9)	73 (70.9)
Length (days) of ICU stay (mean (SD))		11.5 (10.7)	11.415 (10.7)	11.3 (10.8)	12.9 (11.0)
Discharge status					
Recovered at discharge (n, %)		6003 (51.2)	4244 (51.8)	1378 (49.0)	381 (53.1)
Recovered at home (n, %)		3093 (26.4)	2201 (26.9)	722 (25.7)	170 (23.7)
In-hospital death (n, %)		2388 (20.4)	1574 (19.2)	622 (23.6)	152 (21.2)
Transferred (n, %)		201 (1.7)	149 (51.8)	44 (49.0)	8 (53.1)
Unknown (n, %)		32 (0.3)	21 (0.3)	4 (0.1)	7 (1.0)
Length (days) of hospital stay (mean (SD))		12.6 (10.9)	12.1 (10.5)	13.9 (11.7)	12.2 (11.6)

ACEI/ARBs, Angiotensin converting enzyme inhibitors / Angiotensin receptor blockers; ARDS, acute respiratory distress syndrome; ICU, intensive care unit.

1

2

3708

4709

5710

6711

7

8

9

10

11

12

13

14

15

16

17712

18713

19714

20715

21

22

23

24

25

26

27

28

29

30

31

32

33

34

35

36

37

38

39

40

41

42

43

44

45

46

47

48

49

50

51

52

53

54

55

56

57

58


59

60

Table 3. State-arrival extended Cox-Markov multivariate model's transitions hazard ratios (HR, 95%CI) as a function of ACEI/ARBs


Transition		ACEI/ARBs use			
		Model 1	p-value	Model 2	p-value
1	Admission → Severity	1.15 (0.98, 1.36)	0.092	1.10 (0.88, 1.36)	0.409
2	Admission → Recovery	<b>1.07 (1.01, 1.13)</b>	0.027	1.05 (0.98, 1.13)	0.182
3	Admission → Death	<b>0.83 (0.75, 0.93)</b>	0.001	<b>0.80 (0.70, 0.91)</b>	0.001
4	Severity → Recovery	1.16 (0.97, 1.38)	0.098	1.16 (0.93, 1.45)	0.195
5	Severity → Death	0.91 (0.73, 1.13)	0.381	1.11 (0.83, 1.49)	0.485

ACEI/ARBs, Angiotensin converting enzyme inhibitors / Angiotensin receptor blockers Model 1: Adjusted for gender, age (years), prevalent CVD, HBP, DM, and time (days) to severity; Model 2: Further by prevalent obesity, and cognitive issues.



Schematic representation of the competing risk multi-state model and transition event matrix (number (%) patients in each transition)

21x12mm (600 x 600 DPI)



Plots for cumulative transition hazards (A), and state transition probabilities (B), and transition probabilities after transfer to intensive care (C) in a multi-state competing risk model considering ACEI/ARBs use (dashed line) versus no use (solid line).

17x33mm (600 x 600 DPI)

## SUPPLEMENTAL MATERIAL

**Table 1 Supplemental.** Frequency of symptoms and clinical signs compatible with COVID-19 at hospital admission according to ACEI/ARBs use

	Total (n = 11717)	ACEI/ARBs		
		No use (n = 8189, 69.9%)	Use (n = 2810, 23.9%)	Unknown use (n = 718, 6.1%)
<b>Symptoms at admission</b>	11039 (94.2)	7705 (94.1)	2673 (95.1)	661 (92.1)
Fever (n, %)	7187 (61.3)	5112 (62.4)	1653 (58.8)	422 (58.8)
Cough (n, %)	6231 (53.2)	4444 (54.3)	1438 (51.2)	349 (48.6)
Breathlessness (n, %)	5997 (51.2)	4194 (51.2)	1349 (52.0)	376 (47.6)
Weakness (n, %)	4672 (39.9)	3253 (39.7)	1153 (41.0)	266 (37.0)
Pain (n, %)	2622 (22.4)	1931 (23.6)	544 (19.4)	147 (20.5)
Diarrhea (n, %)	1673 (14.3)	1173 (14.3)	438 (15.6)	62 (8.6)
Nausea and vomiting (n, %)	1364 (11.6)	987 (12.1)	311 (11.1)	66 (9.2)
Headache (n, %)	1234 (10.5)	932 (11.4)	245 (8.7)	57 (7.9)
Irritability (n, %)	838 (7.2)	556 (6.8)	243 (8.6)	39 (5.4)
Throat pain (n, %)	700 (6.0)	520 (6.3)	141 (5.0)	39 (5.4)
Anosmia (n, %)	424 (3.6)	334 (4.1)	79 (2.8)	11 (1.5)
Missing (n, %)	855 (7.3)	593 (7.2)	216 (7.7)	46 (6.4)
Runny nose (n, %)	416 (3.6)	301 (3.7)	91 (3.2)	24 (3.3)
<b>Clinical signs at admission</b>	9993 (85.3)	7005 (85.5)	2452 (87.3)	536 (74.7)
Abnormal pulmonary imaging (n, %)*	7396 (63.1)	5271 (64.4)	1835 (65.3)	290 (40.4)
Abnormal pulmonary auscultation (n, %)	5245 (44.8)	3701 (45.2)	1343 (47.8)	201 (28.0)
Dyspnea (n, %)	4966 (42.4)	3462 (42.3)	1196 (42.6)	308 (42.9)
Pharyngitis (n, %)	244 (2.1)	174 (2.1)	58 (2.1)	12 (1.7)
Coma (n, %)	72 (0.6)	50 (0.6)	16 (0.6)	6 (0.8)
Conjunctivitis (n, %)	57 (0.5)	37 (0.5)	18 (0.6)	2 (0.3)
Convulsions (n, %)	15 (0.1)	12 (0.1)	3 (0.1)	0 (0.0)

ACEI; Angiotensin converting enzyme inhibitors, ARBs. Angiotensin receptor blockers

\* Reported as abnormal pulmonary imaging compatible with pneumonia

**Table 2 Supplemental.** Results (OR, 95%CI) of variable selection models of conditions associated with ACEI/ARBs use among factors with COVID-19 prognosis<sup>1</sup>

Variable	Model 1	p-value	Model 2	p-value
Intercept	0.04 (0.03-0.05)	< 0.0001	0.03 (0.03-0.05)	< 0.001
Gender (male)	1.33 (1.21-1.47)	< 0.0001	1.26 (1.12-1.41)	< 0.001
Age	1.01 (1.01-1.02)	< 0.0001	1.01 (1.01-1.02)	< 0.001
CVD	1.71 (1.55-1.90)	< 0.0001	1.65 (1.46-1.87)	< 0.001
Diabetes	1.38 (1.24-1.54)	< 0.0001	1.37 (1.20-1.56)	< 0.001
HBP	5.65 (5.10-6.27)	< 0.0001	5.28 (4.66-6.00)	< 0.001
Obesity			1.33 (1.10-1.59)	< 0.01
Cognitive impairment			0.69 (0.58-0.82)	< 0.001

<sup>1</sup> Backwards stepwise logistic regression with variable selection according to AIC; OR, odds ratio. Model 1 included: Gender, age, HBP, CVD, DM, chronic renal disease, chronic liver disease, chronic lung disease, solid malignant neoplasms, hematological cancers, immunodepression (N=10866); Model 2 included: Model 1 variables plus cognitive impairment, chronic neuro-muscular disease, and obesity, for which a high degree of missingness was observed and represents a complete case analysis (N=7,294)

**Table 3 Supplemental.** State-arrival extended Cox-Markov models transitions hazard ratios (HR, 95%CI) as a function of ACEI/ARBs and identified confounders**Model 1**

Transition	ACEI/ARBs use	Male gender	Age < 70	Age > 70	CVD	HBP	DM	Time to severity
1 Admission to Severity	1.15 (0.98, 1.36) 0.092	<b>1.83 (1.58, 2.12)</b> <0.0001	<b>1.02 (1.01, 1.02)</b> <0.0001	<b>0.90 (0.89, 0.91)</b> <0.0001	0.96 (0.82, 1.12) 0.575	<b>1.35 (1.15, 1.58)</b> <0.0001	0.94 (0.79, 1.11) 0.440	
2 Admission to Recovery	<b>1.07 (1.01, 1.13)</b> 0.027	<b>0.89 (0.85, 0.93)</b> <0.0001	<b>0.97 (0.97, 0.97)</b> <0.0001	<b>0.96 (0.96, 0.97)</b> <0.0001	<b>0.86 (0.81, 0.91)</b> <0.0001	<b>0.95 (0.89, 1.00)</b> 0.043	<b>0.92 (0.87, 0.98)</b> 0.007	
3 Admission to Death	<b>0.83 (0.75, 0.93)</b> 0.001	<b>1.45 (1.31, 1.59)</b> <0.0001	<b>1.11 (1.09, 1.13)</b> <0.0001	<b>1.05 (1.04, 1.05)</b> <0.0001	<b>1.13 (1.02, 1.25)</b> 0.015	1.00 (0.90, 1.10) 0.947	1.09 (0.98, 1.22) 0.102	
4 Severity to recovery	1.16 (0.97, 1.38) 0.098	0.90 (0.77, 1.04) 0.158	<b>0.97 (0.96, 0.97)</b> <0.0001	1.00 (0.98, 1.02) 0.799	1.03 (0.87, 1.21) 0.766	0.95 (0.81, 1.12) 0.570	1.01 (0.85, 1.19) 0.919	<b>0.95 (0.93, 0.97)</b> <0.0001
5 Severity to death	0.91 (0.73, 1.13) 0.381	1.11 (0.91, 1.36) 0.316	<b>1.05 (1.03, 1.06)</b> <0.0001	1.05 (1.03, 1.07) <0.0001	1.07 (0.88, 1.30) 0.494	0.90 (0.74, 1.11) 0.326	<b>1.25 (1.02, 1.53)</b> 0.034	1.02 (1.00, 1.04) 0.056


**Model 2**


Transition	ACEI/ARBs use	Male gender	Age < 70	Age > 70	CVD	HBP	DM	Cognitive issues	Obesity	Time to severity
1 Admission to severity	1.10 (0.88, 1.36) 0.409	<b>1.71 (1.42, 2.08)</b> <0.0001	<b>1.02 (1.01, 1.03)</b> 0.001	<b>0.92 (0.90, 0.93)</b> <0.0001	0.91 (0.74, 1.12) 0.361	<b>1.45 (1.18, 1.79)</b> 0.001	0.96 (0.77, 1.19) 0.686	<b>0.52 (0.35, 0.77)</b> 0.001	<b>1.58 (1.24, 2.02)</b> <0.0001	
2 Admission to recovery	1.05 (0.98, 1.13) 0.182	<b>0.89 (0.84, 0.94)</b> <0.0001	<b>0.97 (0.97, 0.97)</b> <0.0001	<b>0.97 (0.96, 0.97)</b> <0.0001	<b>0.88 (0.82, 0.94)</b> <0.0001	0.97 (0.91, 1.04) 0.377	0.93 (0.87, 1.00) 0.061	<b>0.83 (0.76, 0.91)</b> <0.0001	1.01 (0.92, 1.11) 0.882	
3 Admission to death	<b>0.80 (0.70, 0.91)</b> 0.001	<b>1.48 (1.31, 1.66)</b> <0.0001	<b>1.10 (1.07, 1.12)</b> <0.0001	<b>1.05 (1.04, 1.05)</b> <0.0001	<b>1.13 (1.00, 1.27)</b> 0.048	1.03 (0.91, 1.16) 0.685	1.11 (0.97, 1.27) 0.116	<b>1.36 (1.19, 1.55)</b> <0.0001	0.97 (0.76, 1.24) 0.824	
4 Severity to recovery	1.16 (0.93, 1.45) 0.195	0.90 (0.75, 1.09) 0.272	<b>0.97 (0.96, 0.98)</b> <0.0001	1.00 (0.97, 1.02) 0.861	1.12 (0.91, 1.38) 0.288	1.00 (0.81, 1.24) 0.984	0.95 (0.77, 1.16) 0.597	1.22 (0.84, 1.77) 0.297	0.97 (0.77, 1.22) 0.781	<b>0.94 (0.91, 0.97)</b> <0.0001
5 Severity to death	1.11 (0.83, 1.49) 0.485	1.22 (0.93, 1.60) 0.146	<b>1.04 (1.02, 1.06)</b> <0.0001	<b>1.06 (1.03, 1.08)</b> <0.0001	1.08 (0.82, 1.41) 0.591	0.81 (0.61, 1.08) 0.150	<b>1.44 (1.10, 1.90)</b> 0.008	0.89 (0.53, 1.50) 0.668	0.97 (0.69, 1.36) 0.860	<b>1.04 (1.01, 1.06)</b> 0.003

ACEI/ARBs, Angiotensin converting enzyme inhibitors / Angiotensin receptor blockers; CVD, cardiovascular disease; DM, diabetes mellitus; HBP, high blood pressure. Model 1: Adjusted for gender, age (years), CVD, HBP, DM, and time (days) to severity; Model 2: Further by prevalent obesity, and cognitive issues.



Figure 1 Supplemental.





Stacked probability plot of the state-occupation probabilities in a multi-state competing risk model considering ACEI/ARBs use (A) versus no use (B).

40x24mm (600 x 600 DPI)

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

STROBE Statement—checklist of items that should be included in reports of observational studies

	Item No	Recommendation	Page No
Title and abstract	1	(a) Indicate the study’s design with a commonly used term in the title or the abstract	1
		(b) Provide in the abstract an informative and balanced summary of what was done and what was found	2,3
Introduction			
Background/rationale	2	Explain the scientific background and rationale for the investigation being reported	4
Objectives	3	State specific objectives, including any prespecified hypotheses	5
Methods			
Study design	4	Present key elements of study design early in the paper	5,6
Setting	5	Describe the setting, locations, and relevant dates, including periods of recruitment, exposure, follow-up, and data collection	7,8
Participants	6	(a) Cohort study—Give the eligibility criteria, and the sources and methods of selection of participants. Describe methods of follow-up Case-control study—Give the eligibility criteria, and the sources and methods of case ascertainment and control selection. Give the rationale for the choice of cases and controls Cross-sectional study—Give the eligibility criteria, and the sources and methods of selection of participants	6,7
		(b) Cohort study—For matched studies, give matching criteria and number of exposed and unexposed Case-control study—For matched studies, give matching criteria and the number of controls per case	
Variables	7	Clearly define all outcomes, exposures, predictors, potential confounders, and effect modifiers. Give diagnostic criteria, if applicable	7,8
Data sources/ measurement	8*	For each variable of interest, give sources of data and details of methods of assessment (measurement). Describe comparability of assessment methods if there is more than one group	5,6
Bias	9	Describe any efforts to address potential sources of bias	8-10
Study size	10	Explain how the study size was arrived at	6,7
Quantitative variables	11	Explain how quantitative variables were handled in the analyses. If applicable, describe which groupings were chosen and why	7,8
Statistical methods	12	(a) Describe all statistical methods, including those used to control for confounding	8-10
		(b) Describe any methods used to examine subgroups and interactions	9
		(c) Explain how missing data were addressed	9
		(d) Cohort study—If applicable, explain how loss to follow-up was addressed Case-control study—If applicable, explain how matching of cases and controls was addressed Cross-sectional study—If applicable, describe analytical methods taking account of sampling strategy	9
		(e) Describe any sensitivity analyses	9

Continued on next page

**Results**

Participants	13*	(a) Report numbers of individuals at each stage of study—eg numbers potentially eligible, examined for eligibility, confirmed eligible, included in the study, completing follow-up, and analysed	10, and Fig 1
		(b) Give reasons for non-participation at each stage	10
		(c) Consider use of a flow diagram	Fig 1
Descriptive data	14*	(a) Give characteristics of study participants (eg demographic, clinical, social) and information on exposures and potential confounders	11, Table 1
		(b) Indicate number of participants with missing data for each variable of interest	11, Table 1
		(c) <i>Cohort study</i> —Summarise follow-up time (eg, average and total amount)	11, Table 2
Outcome data	15*	<i>Cohort study</i> —Report numbers of outcome events or summary measures over time	10-11, table 2
		<i>Case-control study</i> —Report numbers in each exposure category, or summary measures of exposure	
		<i>Cross-sectional study</i> —Report numbers of outcome events or summary measures	
Main results	16	(a) Give unadjusted estimates and, if applicable, confounder-adjusted estimates and their precision (eg, 95% confidence interval). Make clear which confounders were adjusted for and why they were included	12 table 3
		(b) Report category boundaries when continuous variables were categorized	All tables
		(c) If relevant, consider translating estimates of relative risk into absolute risk for a meaningful time period	
Other analyses	17	Report other analyses done—eg analyses of subgroups and interactions, and sensitivity analyses	13, Table 3

**Discussion**

Key results	18	Summarise key results with reference to study objectives	13-16
Limitations	19	Discuss limitations of the study, taking into account sources of potential bias or imprecision. Discuss both direction and magnitude of any potential bias	16-18
Interpretation	20	Give a cautious overall interpretation of results considering objectives, limitations, multiplicity of analyses, results from similar studies, and other relevant evidence	14-16
Generalisability	21	Discuss the generalisability (external validity) of the study results	14-16

**Other information**

Funding	22	Give the source of funding and the role of the funders for the present study and, if applicable, for the original study on which the present article is based	19
---------	----	---	----

\*Give information separately for cases and controls in case-control studies and, if applicable, for exposed and unexposed groups in cohort and cross-sectional studies.

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

**Note:** An Explanation and Elaboration article discusses each checklist item and gives methodological background and published examples of transparent reporting. The STROBE checklist is best used in conjunction with this article (freely available on the Web sites of PLoS Medicine at <http://www.plosmedicine.org/>, Annals of Internal Medicine at <http://www.annals.org/>, and Epidemiology at <http://www.epidem.com/>). Information on the STROBE Initiative is available at [www.strobe-statement.org](http://www.strobe-statement.org).

For peer review only